







INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07C 275/24, A61K 31/17 C07D 215/48, C07C 317/44 C07C 275/22, A61K 31/47	A1	(11) International Publication Number: WO 93/23368 (43) International Publication Date: 25 November 1993 (25.11.93)
(21) International Application Number: PCT/US93/04806 (22) International Filing Date: 20 May 1993 (20.05.93) (30) Priority data: 07/886,558 20 May 1992 (20.05.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/886,558 (CON) Filed on 20 May 1992 (20.05.92) (71) Applicants (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Department, P.O. Box 5110, Chicago, IL 60680-5110 (US). THE MONSANTO COMPANY [US/US]; 800 North Lindbergh Boulevard, St. Louis, MO 63167 (US). (72) Inventors; and (75) Inventors/Applicants (for US only) : TALLEY, John, J. [US/US]; 1079 Darwick Court, St. Louis, MO 63132 (US). GETMAN, Daniel, P. [US/US]; 66 Sunnyhill Court, Chesterfield, MO 63017 (US). DECRESCENZO, Gary, A. [US/US]; 536 Schrader Farm Drive, St. Peters, MO 63376 (US). LIN, Ko-Chung [CN/US]; 12790 Polo Parc Drive, St. Louis, MO 63146 (US). VAZQUEZ, Michael, L. [US/US]; 233 Saratoga Court, Gurnee, IL 60031 (US). MUELLER, Richard, A. [US/US]; 562 Stonegate Terrace, Glencoe, IL 60022 (US). REED, Kathryn, L. [US/US]; 7301-104 Brookmont Drive, Raleigh, NC 27163		(US) HEINTZ, Robert, M. [US/US]; 603 Nancy Place, Ballwin, MO 63021 (US). CLARE, Michael [GB/US]; 5154 West Brown Street, Skokie, IL 60077 (US). FRESKOS, John, N. [US/US]; 7572 York, Clayton, MO 63105 (US). SUN, Eric, T. [US/US]; 7614 Palmilla Drive, Apartment 58, San Diego, CA 92122 (US). (74) Agents: KEANE, J., Timothy et al.; G.D. Searle & Co., Corporate Patent Department, P.O. Box 5110, Chicago, IL 60680-5110 (US). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: UREA-CONTAINING HYDROXYETHYLAMINE COMPOUNDS AS RETROVIRAL PROTEASE INHIBITORS

**(57) Abstract**

Compounds represented by formula (I) or a pharmaceutically acceptable salt, prodrug or ester thereof, wherein A represents radicals represented by formulae (A1, A2, A3) (values for the variables given herein), are effective as retroviral protease inhibitors, and in particular as inhibitors of HIV protease.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

UREA-CONTAINING HYDROXYETHYLAMINE COMPOUNDS AS RETROVIRAL
PROTEASE INHIBITORS

This application is a continuation-in-part of Patent Application Serial No. PCT/US91/8613, filed November 18, 1991, which is a continuation-in-part of 07/789,646, filed November 14, 1991, which is a continuation-in-part of U.S. Patent Application Serial No. 07/615,210, filed November 19, 1990.

10

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to retroviral protease inhibitors and, more particularly, relates to novel compounds and a composition and method for inhibiting retroviral proteases. This invention, in particular, relates to urea-containing hydroxyethylamine protease inhibitor compounds, a composition and method for inhibiting retroviral proteases such as human immunodeficiency virus (HIV) protease and for treatment or prophylaxis a retroviral infection, e.g., an HIV infection. The subject invention also relates to processes for making such compounds as well as to intermediates useful in such processes.

25

2. Related Art

During the replication cycle of retroviruses, gag and gag-pol gene products are translated as proteins. These proteins are subsequently processed by a virally encoded protease (or proteinase) to yield viral enzymes and structural proteins of the virus core. Most commonly, the gag precursor proteins are processed into the core proteins and the pol precursor proteins are processed into the viral enzymes, e.g., reverse transcriptase and retroviral protease. It has been shown that correct processing of the precursor proteins by the retroviral protease is necessary for assembly of infectious virions. For example, it has been shown that frameshift mutations in the protease region of

the pol gene of HIV prevents processing of the gag precursor protein. It has also been shown through site-directed mutagenesis of an aspartic acid residue in the HIV protease that processing of the gag precursor protein is prevented.

5 Thus, attempts have been made to inhibit viral replication by inhibiting the action of retroviral proteases.

Retroviral protease inhibition typically involves a transition-state mimetic whereby the retroviral protease

10 is exposed to a mimetic compound which binds (typically in a reversible manner) to the enzyme in competition with the gag and gag-pol proteins to thereby inhibit replication of structural proteins and, more importantly, the retroviral protease itself. In this manner, retroviral replication

15 proteases can be effectively inhibited.

Several classes of mimetic compounds have been proposed, particularly for inhibition of proteases, such as for inhibition of HIV protease. Such mimetics include

20 hydroxyethylamine isosteres and reduced amide isosteres. See, for example, EP O 346 847; EP O 342,541; Roberts et al, "Rational Design of Peptide-Based Proteinase Inhibitors, "Science, 248, 358 (1990); and Erickson et al, "Design Activity, and 2.8Å Crystal Structure of a C₂ Symmetric

25 Inhibitor Complexed to HIV-1 Protease," Science, 249, 527 (1990).

Several classes of mimetic compounds are known to be useful as inhibitors of the proteolytic enzyme renin. See, for example, U.S. No. 4,599,198; U.K. 2,184,730; G.B. 2,209,752; EP O 264 795; G.B. 2,200,115 and U.S. SIR H725. 5 Of these, G.B. 2,200,115, GB 2,209,752, EP O 264,795, U.S. SIR H725 and U.S. 4,599,198 disclose urea-containing hydroxyethylamine renin inhibitors. However, it is known that, although renin and HIV proteases are both classified as aspartyl proteases, compounds which are effective renin 10 inhibitors generally cannot be predicted to be effective HIV protease inhibitors.

BRIEF DESCRIPTION OF THE INVENTION


15

The present invention is directed to virus inhibiting compounds and compositions. More particularly, the present invention is directed to retroviral protease inhibiting compounds and compositions, to a method of 20 inhibiting retroviral proteases, to processes for preparing the compounds and to intermediates useful in such processes. The subject compounds are characterized as urea-containing hydroxyethylamine inhibitor compounds.

25

DETAILED DESCRIPTION OF THE INVENTION


In accordance with the present invention, there is provided a retroviral protease inhibiting compound of the formula:




(Formula I)

or a pharmaceutically acceptable salt, prodrug or ester thereof wherein:


A represents radicals represented by the formulas:



(A1)



(A2)



(A3)

- wherein R represents hydrogen, alkoxycarbonyl, aralkoxycarbonyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkoxycarbonyl, cycloalkylalkanoyl, alkanoyl, aralkanoyl, aroyl, aryloxy carbonyl, aryloxy carbonylalkyl, aryloxyalkanoyl, heterocyclylcarbonyl, heterocyclyloxy carbonyl, heterocyclylalkanoyl, heterocyclylalkoxycarbonyl, heteroaralkoxycarbonyl, heteroaryloxy carbonyl, heteroaroyl, heteroaralkanoyl, alkyl, aryl, aralkyl, aryloxyalkyl, heteroaryloxyalkyl,

hydroxyalkyl, aminocarbonyl, aminoalkanoyl, and mono- and disubstituted aminocarbonyl and mono- and disubstituted aminoalkanoyl radicals wherein the substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkanoyl radical, said substituents along with the nitrogen atom to which they are attached form a heterocyclyl or heteroaryl radical;

R' represents hydrogen and radicals as defined for R³ or R and R' together with the nitrogen to which they are attached form a heterocycloalkyl or heteroaryl radical, or when A is Al, R' represents hydrogen, radicals as defined for R³ and arakloxy carbonylalkyl and aminocarbonylalkyl and aminocarbonyl radicals wherein said amino group may be mono- or disubstituted with substituents selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl and heterocycloalkylalkyl radicals;

t represents either 0 or 1;

R¹ represents hydrogen, -CH₂SO₂NH₂, -CO₂CH₃, -CONHCH₃, -CON(CH₃)₂, -CH₂C(O)NHCH₃, -CH₂C(O)N(CH₃)₂, -CONH₂, -C(CH₃)₂(SCH₃), -C(CH₃)₂(S[O]CH₃), -C(CH₃)₂(S[O]CH₃), alkyl, haloalkyl, alkenyl, alkynyl and cycloalkyl radicals and amino acid side chains selected from asparagine, S-methyl cysteine and the corresponding sulfoxide and sulfone derivatives thereof, glycine, leucine, isoleucine, allo-isoleucine, tert-leucine, phenylalanine, ornithine, alanine, histidine, norleucine, glutamine, valine, threonine, serine, aspartic acid, beta-cyano alanine, and allothreonine side chains;

R^{1'} and R^{1''} independently represent hydrogen and radicals as defined for R¹, or one of R^{1'} and R^{1''}; together with R¹ and the carbon atoms to which R¹, R^{1'} and R^{1''} are attached, represent a cycloalkyl radical;

R² represents alkyl, aryl, cycloalkyl, cycloalkylalkyl and aralkyl radicals, which radicals are optionally substituted with a group selected from -NO₂, -OR⁹, -SR⁹, and halogen radicals, wherein R⁹ represents hydrogen and alkyl radicals;

R³ represents alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, heterocycloalkylalkyl, aryl, aralkyl, heteroaralkyl, aminoalkyl and mono- and disubstituted aminoalkyl radicals, wherein said substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkyl radical, said substituents along with the nitrogen atom to which they are attached, form a heterocycloalkyl or a heteroaryl radical;

X' represents O, C(R¹⁷) where R¹⁷ represents hydrogen and alkyl radicals, and N;

Y, Y', and Y'' independently represent O, S and NR¹⁵ wherein R¹⁵ represents hydrogen and radicals as defined for R³;

R⁴ and R⁵ independently represent hydrogen and radicals as defined by R³, or R⁴ and R⁵ together with the nitrogen atom to which they are bonded represent heterocycloalkyl and heteroaryl radicals;


R⁶ represents hydrogen and alkyl radicals;

R²⁰, R²¹, R³⁰, R³¹ and R³² represent radicals as defined for R¹, or one of R¹ and R³⁰ together with one of R³¹ and R³²

and the carbon atoms to which they are attached form a cycloalkyl radical; and

- 5 R^{33} and R^{34} independently represent hydrogen and radicals as defined for R^3 , or R^{33} and R^{34} together with X' represent cycloalkyl, aryl, heterocyclyl and heteroaryl radicals, provided that when X' is O, R^{34} is absent.

- 10 A preferred class of retroviral inhibitor compounds of the present invention are those represented by the formula:



- 15 or a pharmaceutically acceptable salt, prodrug or ester thereof, preferably wherein the stereochemistry about the hydroxy group is designated as (R);

- 20 R represents hydrogen, alkoxycarbonyl, aralkoxycarbonyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkoxycarbonyl, cycloalkylalkanoyl, alkanoyl, aralkanoyl, aroyl, aryloxy carbonyl, aryloxy carbonylalkyl, aryloxyalkanoyl, heterocyclylcarbonyl, heterocyclylalkoxycarbonyl, heterocyclylalkanoyl, heterocyclylalkoxycarbonyl, heteroaralkanoyl, heteroaralkoxycarbonyl, heteroaryloxy carbonyl, heteroaroyl, alkyl, alkenyl, cycloalkyl, aryl, aralkyl, aryloxyalkyl, heteroaryloxyalkyl, hydroxyalkyl, aminocarbonyl, aminoalkanoyl, and mono- and disubstituted aminocarbonyl and mono- and disubstituted aminoalkanoyl radicals wherein the substituents are selected from alkyl, 30 aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, heterocycloalkylalkyl radicals, or where said aminoalkanoyl radical is

disubstituted, said substituents along with the nitrogen atom to which they are attached form a heterocycloalkyl or heteroaryl radical;

- 5 R' represents hydrogen and radicals as defined for R³ or R and R' together with the nitrogen to which they are attached represent heterocycloalkyl and heteroaryl radical;

- R¹ represents hydrogen, -CH₂SO₂NH₂, -CH₂CO₂CH₃, -CO₂CH₃,
10 -CONH₂, -CHC(O)NHCH₃, -C(CH₃)₂(SCH₃), -C(CH₃)₂(S[O]CH₃),
-C(CH₃)₂(S[O]₂CH₃), alkyl, haloalkyl, alkenyl, alkynyl and cycloalkyl radicals, and amino acid side chains selected from asparagine, S-methyl cysteine and the sulfoxide (SO) and sulfone (SO₂) derivatives thereof, isoleucine,
15 allo-isoleucine, alanine, leucine, tert-leucine, phenylalanine, ornithine, histidine, norleucine, glutamine, threonine, glycine, allo-threonine, serine, aspartic acid, beta-cyano alanine and valine side chains;

- 20 R^{1'} and R^{1''} independently represent hydrogen and radicals as defined for R¹, or one of R^{1'} and R^{1''}, together with R¹ and the carbon atoms to which R¹, R^{1'} and R^{1''} are attached, represent a cycloalkyl radical;

- 25 R² represents alkyl, aryl, cycloalkyl, cycloalkylalkyl, and aralkyl radicals, which radicals are optionally substituted with a group selected from alkyl radicals, NO₂, OR⁹ and SR⁹ wherein R⁹ represents hydrogen and alkyl radicals, and halogen radicals;

- 30 R³ represents alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, heterocycloalkylalkyl, aryl, aralkyl, heteroaralkyl, aminoalkyl and mono- and disubstituted
35 aminoalkyl radicals, wherein said substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, and

heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkyl radical, said substituents along with the nitrogen atom to which they are attached, form a heterocycloalkyl or a heteroaryl radical;

5

R⁴ and R⁵ independently represent hydrogen and radicals as defined by R³, or together with a nitrogen atom to which they are bonded represent heterocycloalkyl and heteroaryl radicals;

10

t represents 0 or 1;

Y and Y' independently represent O, S, and NR¹⁵ wherein R¹⁵ represents hydrogen and radicals as defined for R³.


15

Preferably, Y and Y' represent O.

Preferably, R³ represents radicals as defined above which contain no α -branching, e.g., as in an isopropyl radical or a t-butyl radical. The preferred radicals are those which contain a -CH₂- moiety between the nitrogen of the urea and the remaining portion of the radical. Such preferred groups include, but are not limited to, benzyl, isobutyl, n-butyl, isoamyl, cyclohexylmethyl and the like.

25

Another preferred class of compounds are those represented by the formula:



(Formula III)


30

or a pharmaceutically acceptable salt, prodrug or ester thereof wherein t, Y, Y', R', R¹, R², R³, R⁴, R⁵, R²⁰ and R²¹ are as defined above with respect to Formula I.

35

Preferably, the stereochemistry about the hydroxyl group is designated as R.

Yet another preferred class of compounds are those represented by the formula:



(Formula IV)

or a pharmaceutically acceptable salt, prodrug or ester thereof, preferably wherein the stereochemistry about the hydroxy group is designated as R, wherein t, X', Y, Y', Y", R¹, R², R³, R⁴, R⁵, R³⁰, R³¹, R³², R³³ and R³⁴ are as defined above with respect to Formula I.

As utilized herein, the term "alkyl", alone or in combination, means a straight-chain or branched-chain alkyl radical containing from 1 to about 10, preferably from 1 to about 8, carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl and the like. The term "alkenyl", alone or in combination, means a straight-chain or branched-chain hydrocarbon radical having one or more double bonds and containing from 2 to about 18 carbon atoms preferably from 2 to about 8 carbon atoms. Examples of suitable alkenyl radicals include ethenyl, propenyl, alkyl, 1,4-butadienyl and the like. The term "alkynyl", alone or in combination, means a straight-chain hydrocarbon radical having one or more triple bonds and containing from 2 to about 10 carbon atoms. Examples of alkynyl radicals include ethynyl, propynyl, (propargyl), butynyl and the like. The term "alkoxy", alone or in combination, means an alkyl ether radical wherein the term alkyl is as defined above. Examples of suitable alkyl ether radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy and the like.

The term "cycloalkyl", alone or in combination, means an alkyl radical which contains from about 3 to about 8 carbon atoms and is cyclic. The term "cycloalkylalkyl" means an alkyl radical as defined above which is substituted by a cycloalkyl radical containing from about 3 to about 8, preferably from about 3 to about 6, carbon atoms. Examples of such cycloalkyl radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like. The term "aryl", alone or in combination, means a phenyl or naphthyl radical which optionally carries one or more substituents selected from alkyl, alkoxy, halogen, hydroxy, amino, nitro and the like, such as phenyl, p-tolyl, 4-methoxyphenyl, 4-(tert-butoxy)phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-hydroxyphenyl, 1-naphthyl, 2-naphthyl, and the like. The term "aralkyl", alone or in combination, means an alkyl radical as defined above in which one hydrogen atom is replaced by an aryl radical as defined above, such as benzyl, 2-phenylethyl and the like. The term "aralkoxy carbonyl", alone or in combination, means a radical of the formula $-C(O)-O-$ aralkyl in which the term "aralkyl" has the significance given above. An example of an aralkoxycarbonyl radical is benzyloxycarbonyl. The term "aryloxy" means a radical of the formula aryl- $O-$ in which the term aryl has the significance given above. The term "alkanoyl", alone or in combination, means an acyl radical derived from an alkanecarboxylic acid, examples of which include acetyl, propionyl, butyryl, valeryl, 4-methylvaleryl, and the like. The term "cycloalkylcarbonyl" means an acyl group derived from a monocyclic or bridged cycloalkanecarboxylic acid such as cyclopropanecarbonyl, cyclohexanecarbonyl, adamantanecarbonyl, and the like, or from a benz-fused monocyclic cycloalkanecarboxylic acid which is optionally substituted by, for example, alkanoylamino, such as 1,2,3,4-tetrahydro-2-naphthoyl, 2-acetamido-1,2,3,4-tetrahydro-2-naphthoyl. The term "aralkanoyl" means an acyl radical derived from an aryl-substituted alkanecarboxylic acid such as phenylacetyl, 3-phenylpropionyl (hydrocinnamoyl),

4-phenylbutyryl, (2-naphthyl)acetyl, 4-chlorohydrocinnamoyl, 4-aminohydrocinnamoyl, 4-methoxyhydrocinnamoyl, and the like. The term "aroyl" means an acyl radical derived from an aromatic carboxylic acid. Examples of such radicals include aromatic carboxylic acids, an optionally substituted benzoic or naphthoic acid such as benzoyl, 4-chlorobenzoyl, 4-carboxybenzoyl, 4-(benzyloxycarbonyl)benzoyl, 1-naphthoyl, 2-naphthoyl, 6-carboxy-2-naphthoyl, 6-(benzyloxycarbonyl)-2-naphthoyl, 3-benzyloxy-2-naphthoyl, 3-hydroxy-2-naphthoyl, 3-(benzyloxyformamido)-2-naphthoyl, and the like. The heterocyclyl or heterocycloalkyl portion of a heterocyclylcarbonyl, heterocyclilyloxycarbonyl, heterocyclylalkoxycarbonyl, or heterocyclialkyl group or the like is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic heterocycle which contains one or more hetero atoms selected from nitrogen, oxygen and sulphur, which is optionally substituted on one or more carbon atoms by halogen, alkyl, alkoxy, oxo, and the like, and/or on a secondary nitrogen atom (i.e., -NH-) by alkyl, aralkoxycarbonyl, alkanoyl, phenyl or phenylalkyl or on a tertiary nitrogen atom (i.e. = N-) by oxido and which is attached via a carbon atom. The heteroaryl portion of a heteroaroyl, heteroaryloxycarbonyl, or a heteroaralkoxy carbonyl group or the like is an aromatic monocyclic, bicyclic, or tricyclic heterocycle which contains the hetero atoms and is optionally substituted as defined above with respect to the definition of heterocyclyl. Examples of such heterocyclyl and heteroaryl groups are pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiamorpholinyl, pyrrolyl, imidazolyl (e.g., imidazol-4-yl, 1-benzyloxycarbonylimidazol-4-yl, etc.), pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, furyl, thienyl, triazolyl, oxazolyl, thiazolyl, indolyl (e.g., 2-indolyl, etc.), quinolinyl, (e.g., 2-quinolinyl, 3-quinolinyl, 1-oxido-2-quinolinyl, etc.), isoquinolinyl (e.g., 1-isoquinolinyl, 3-isoquinolinyl, etc.), tetrahydroquinolinyl (e.g., 1,2,3,4-tetrahydro-2-quinolyl, etc.),


1,2,3,4-tetrahydroisoquinolinyl (e.g., 1,2,3,4-tetrahydro-1-oxo-isoquinolinyl, etc.), quinoxalinyl, β -carbolinyl, 2-benzofurancarbonyl, 1-,2-,4- or 5-benzimidazolyl, and the like. The term "cycloalkylalkoxycarbonyl" means an acyl group derived from a cycloalkylalkoxycarboxylic acid of the formula cycloalkylalkyl-O-COOH wherein cycloalkylalkyl has the significance given above. The term "aryloxyalkanoyl" means an acyl radical of the formula aryl-O-alkanoyl wherein aryl and alkanoyl have the significance given above. The term "heterocyclyloxycarbonyl" means an acyl group derived from heterocyclyl-O-COOH wherein heterocyclyl is as defined above. The term "heterocyclylalkanoyl" is an acyl radical derived from a heterocyclyl-substituted alkane carboxylic acid wherein heterocyclyl has the significance given above. The term "heterocyclylalkoxycarbonyl" means an acyl radical derived from a heterocyclyl-substituted alkane-O-COOH wherein heterocyclyl has the significance given above. The term "heteroaryloxycarbonyl" means an acyl radical derived from a carboxylic acid represented by heteroaryl-O-COOH wherein heteroaryl has the significance given above. The term "aminocarbonyl" alone or in combination, means an amino-substituted carbonyl (carbamoyl) group derived from an amino-substituted carboxylic acid wherein the amino group can be a primary, secondary or tertiary amino group containing substituents selected from hydrogen, and alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl radicals and the like. The term "aminoalkanoyl" means an acyl group derived from an amino-substituted alkanecarboxylic acid wherein the amino group can be a primary, secondary or tertiary amino group containing substituents selected from hydrogen, and alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl radicals and the like. The term "halogen" means fluorine, chlorine, bromine or iodine. The term "haloalkyl" means an alkyl radical having the significance as defined above wherein one or more hydrogens are replaced with a halogen. Examples of such haloalkyl radicals include chloromethyl, I-bromoethyl, fluoromethyl, difluoromethyl, trifluoromethyl,

1,1,1-trifluoroethyl and the like. The term "leaving group" generally refers to groups readily displaceable by a nucleophile, such as an amine, a thiol or an alcohol nucleophile. Such leaving groups are well known and include
5 carboxylates, N-hydroxysuccinimide, N-hydroxybenzotriazole, halides, triflates, tosylates -OR and -SR and the like. Preferred leaving groups are indicated herein where appropriate.

10 Procedures for preparing the compounds of Formula I are set forth below. It should be noted that the general procedure is shown as it relates to preparation of compounds having the specified stereochemistry, for example, wherein
15 the stereochemistry about the hydroxyl group is designated as (R). However, such procedures are generally applicable, as illustrated in Example 45, to those compounds of opposite configuration, e.g., where the stereochemistry about the hydroxyl group is (S).

20 Preparation of Compounds of Formula II


The compounds of the present invention represented by Formula II above can be prepared utilizing the following general procedure. An N-protected chloroketone derivative
25 of an amino acid having the formula:



wherein P represents an amino protecting group, and R² is as
30 defined above, is reduced to the corresponding alcohol utilizing an appropriate reducing agent. Suitable amino protecting groups are well known in the art and include carbobenzoxy, butyryl, t-butoxycarbonyl, acetyl, benzoyl and the like. A preferred amino protecting group is

carbobenzoxy. A preferred N-protected chloroketone is N-benzyloxycarbonyl-L-phenylalanine chloromethyl ketone. A preferred reducing agent is sodium borohydride. The reduction reaction is conducted at a temperature of from
5 -10°C to about 25°C, preferably at about 0°C, in a suitable solvent system such as, for example, tetrahydrofuran, and the like. The N-protected chloroketones are commercially available from Bachem, Inc., Torrance, California. Alternatively, the chloroketones can be prepared by the
10 procedure set forth in S. J. Fittkau, J. Prakt. Chem., 315, 1037 (1973), and subsequently N-protected utilizing procedures which are well known in the art.


The resulting alcohol is then reacted, preferably
15 at room temperature, with a suitable base in a suitable solvent system to produce an N-protected amino epoxide of the formula:



20 wherein P and R² are as defined above. Suitable solvent systems for preparing the amino epoxide include ethanol, methanol, isopropanol, tetrahydrofuran, dioxane, and the like including mixtures thereof. Suitable bases for
25 producing the epoxide from the reduced chloroketone include potassium hydroxide, sodium hydroxide, potassium t-butoxide, DBU and the like. A preferred base is potassium hydroxide.


Alternatively, a protected amino epoxide can be
30 prepared starting with an L-amino acid which is reacted with a suitable amino-protecting group in a suitable solvent to produce an amino-protected L-amino acid ester of the formula:

16



wherein P, P¹ and P² independently represent hydrogen and amino-protecting groups as defined above with respect to P, provided that P¹ and P² are not both hydrogen; and R² is as defined above.


The amino-protected L-amino acid ester is then reduced, to the corresponding alcohol. For example, the amino-protected L-amino acid ester can be reduced with diisobutylaluminum hydride at -78° C in a suitable solvent such as toluene. The resulting alcohol is then converted, by way of a Swern Oxidation, to the corresponding aldehyde of the formula:



wherein P¹, P² and R² are as defined above. Thus, a dichloromethane solution of the alcohol is added to a cooled (-75 to -68° C) solution of oxalyl chloride in dichloromethane and DMSO in dichloromethane and stirred for 35 minutes.

The aldehyde resulting from the Swern Oxidation is then reacted with a halomethyl lithium reagent, which reagent is generated in situ by reacting an alkyl lithium or aryllithium compound with a dihalomethane represented by the formula X¹CH₂X² wherein X¹ and X² independently represent I, Br or Cl. For example, a solution of the aldehyde and chloriodomethane in THF is cooled to -78° C and a solution of n-butyllithium in hexane is added. The resulting

product is a mixture of diastereomers of the corresponding amino-protected epoxides of the formulas:



5

The diastereomers can be separated by chromatography or, alternatively, once reacted in subsequent steps the diastereomeric products can be separated.

10

The amino epoxide is then reacted, in a suitable solvent system, with an equal amount, or preferably an excess of, a desired amine of the formula:




15

wherein R^3 is hydrogen or is as defined above. The reaction can be conducted over a wide range of temperatures, e.g., from about 10°C to about 100°C, but is preferably, but not necessarily, conducted at a temperature at which the solvent begins to reflux. Suitable solvent systems include those wherein the solvent is an alcohol, such as methanol, ethanol, isopropanol, and the like, ethers such as tetrahydrofuran, dioxane and the like, and toluene, N,N-dimethylformamide, dimethyl sulfoxide, and mixtures thereof. A preferred solvent is isopropanol. Exemplary amines corresponding to the formula R^3NH_2 include benzyl amine, isobutylamine, n-butyl amine, isopentyl amine, isoamylamine, cyclohexanemethyl amine, naphthylene methyl amine and the like. The resulting product is a 3-(N-protected amino)-3-(R^2)-1-(NHR^3)-propan-2-ol derivative


30

(hereinafter referred to as an amino alcohol) can be represented by the formula:




wherein P, R² and R³ are as described above.

For producing compounds of Formula II wherein R⁵ is hydrogen, the resulting amino alcohol described above is then reacted, in a suitable solvent system, with an isocyanate of the formula R⁴NCO wherein R⁴ is as defined above. Where Y in Formula I above is sulfur, the resulting amino alcohol is reacted with an isothiocyanate of the formula R⁴NCS under similar conditions. Suitable solvent systems include tetrahydrofuran, methylene chloride, and the like and mixtures thereof. The resulting product is a urea derivative of the amino alcohol or the corresponding sulfur analog thereof and can be represented by the formula:




wherein P, Y, R², R³ and R⁴ are as defined above. The isocyanates of the formula R⁴NCO can be prepared by the reaction of an amine (R³NH₂) with phosgene, triphosgene, carbodiimidazole, or carbonate ((RO)₂CO) under conditions well-known in the art. The isothiocyanates of the formula R⁴NCS can be prepared by similar procedures, e.g., reaction of the amine with thiophosgene, which are also well known in the art. In addition, the isocyanates and isothiocyanates are commercially available from Aldrich Chemical Company.

For preparing compounds of Formula II wherein R⁵ is other than hydrogen, the resulting amino alcohol described above is then reacted, in a suitable solvent system, with a compound represented by the formula:



wherein R⁴ and R⁵ are as described above and L represents a leaving group such as a halide, e.g., chloride, imidazole radical, the radical p-NO₂-(C₆H₄)-O-, and the like. A preferred compound represented by this formula is a carbamoyl chloride. The corresponding sulfur analogs can be utilized where Y of Formula II is S.


The urea derivative of the amino alcohol and the corresponding sulfur analog can be represented by the formula:



Following preparation of the urea derivative, or corresponding analogs wherein Y is S, the amino protecting group P is removed under conditions which will not affect the remaining portion of the molecule. These methods are well known in the art and include acid hydrolysis, hydrogenolysis and the like. A preferred method involves removal of the protecting group, e.g., removal of a carbobenzoxy group, by hydrogenolysis utilizing palladium on carbon in a suitable solvent system such as an alcohol, acetic acid, and the like or mixtures thereof. Where the protecting group is a t-butoxycarbonyl group, it can be removed utilizing an inorganic or organic acid, e.g., HCl or trifluoroacetic acid, in a suitable solvent system, e.g.,

dioxane or methylene chloride. The resulting product is the amine salt derivative. Following neutralization of the salt, the amine is then reacted with an amino acid or corresponding derivative thereof represented by the formula

5 (PN[CR^{1'} R^{1''}]^t CH(R¹)COOH) wherein t, R¹, R^{1'} and R^{1''} are as defined above, to produce the antiviral compounds of the present invention having the formula:




10

wherein t, P, R¹, R^{1'}, R^{1''}, R², R³, R⁴, R⁵ and Y are as defined above. Preferred protecting groups in this instance are a benzyloxycarbonyl group or a t-butoxycarbonyl group. Where the amine is reacted with a derivative of an amino acid, e.g., when t=1 and R^{1'} and R^{1''} are both H, so that the amino acid is a β-amino acid, such β-amino acids can be prepared according to the procedure set forth in a copending application, U. S. Serial No. 07/345,808. Where t is 1, one of R^{1'} and R^{1''} is H and R¹ is hydrogen so that the amino acid is a homo-β-amino acid, such homo-β-amino acids can be prepared by the procedure set forth in a copending application, U.S. Serial No. 07/853,561. Where t is 0 and R¹ is alkyl, cycloalkyl, -CH₂SO₂NH₂ or an amino acid side chain, such materials are well known and many are

25 commercially available from Sigma-Aldrich.

The N-protecting group can be subsequently removed, if desired, utilizing the procedures described above, and then reacted with a carboxylate represented by

30 the formula:




wherein R is as defined above and L is an appropriate


leaving group such as a halide. Preferably, where R¹ is a side chain of a naturally occurring α-amino acid, R is a 2-quinoline carbonyl group derived from N-hydroxysuccinimide-2-quinoline carboxylate, i.e., L is hydroxy succinimide. A solution of the free amine (or amine acetate salt) and about 1.0 equivalent of the carboxylate are mixed in an appropriate solvent system and optionally treated with up to five equivalents of a base such as, for example, N-methylmorpholine, at about room temperature. Appropriate solvent systems include tetrahydrofuran, methylene chloride or N,N-dimethylformamide, and the like, including mixtures thereof.

Preparation of Compounds of Formula III

A mercaptan of the formula RSH is reacted with a substituted methacrylate of the formula:




by way of a Michael Addition. The Michael Addition is conducted in a suitable solvent and in the presence of a suitable base, to produce the corresponding thiol derivative represented by the formula:



wherein R' and R¹ represent radicals defined above; R²⁰ and R²¹ represent hydrogen and radicals as defined for R¹; and R²² represents radicals as defined by R³. Suitable solvents


in which the Michael Addition can be conducted include alcohols such as, for example, methanol, ethanol, butanol and the like, as well as ethers, e.g., THF, and acetonitrile, DMF, DMSO, and the like, including mixtures thereof. Suitable bases include Group I metal alkoxides such as, for example sodium methoxide, sodium ethoxide, sodium butoxide and the like as well as Group I metal hydrides, such as sodium hydride, including mixtures thereof.

The thiol derivative is converted into the corresponding sulfone of the formula:



by oxidizing the thiol derivative with a suitable oxidation agent in a suitable solvent. Suitable oxidation agents include, for example, hydrogen peroxide, sodium meta-perborate, oxone (potassium peroxy monosulfate), meta-chloroperoxybenzoic acid, and the like, including mixtures thereof. Suitable solvents include acetic acid (for sodium meta-perborate) and, for other peracids, ethers such as THF and dioxane, and acetonitrile, DMF and the like, including mixtures thereof.

The sulfone is then converted to the corresponding free acid of the formula:



5

utilizing a suitable base, e.g., lithium hydroxide, sodium hydroxide, and the like, including mixtures thereof, in a suitable solvent, such as, for example, THF, acetonitrile, DMF, DMSO, methylene chloride and the like, including mixtures thereof.


10

The free acid is then coupled, utilizing, as described above, procedures well known in the art, to the urea derivative, or analog thereof, of an amino alcohol which is described above for the preparation of compounds of Formula II. The resulting product is a compound represented by Formula III.

15

Alternatively, one can couple the urea isostere to the commercially available acid,


20




remove the thioacetyl group with a suitable base, such as hydroxide, or an amine, such as ammonia, and then react the resulting thiol with an alkylating agent, such as an alkyl halide, tosylate or mesylate to afford compounds at the following structure:

25


24




The sulfur can then be oxidized to the corresponding sulfone using suitable oxidizing agents, as described above, to afford the desired compounds of the following structure:



Alternatively, to prepare compounds of Formula III, a substituted methacrylate of the formula:




wherein L represents a leaving group as previously defined, R^{35} and R^{36} represent hydrogen and radicals as defined for R^1 ; and R^{37} represents alkyl, aralkyl, cycloalkyl and cycloalkylalkyl radicals, is reacted with a suitable sulfonating agent, such as, for example, a sulfinic acid represented by the formula RSO_2M , wherein R represents radicals as defined above and M represents a metal adapted to form a salt of the acid, e.g., sodium, to produce the corresponding sulfone represented by the formula:




25

wherein R', R³⁵, R³⁶ and R³⁷ are as defined above. The sulfone is then hydrolyzed in the presence of a suitable base, such as lithium hydroxide, sodium hydroxide and the like, to the compound represented by the formula:




wherein R', R³⁵ and R³⁶ represent radicals as defined above. The resulting compound is then asymmetrically hydrogenated utilizing an asymmetric hydrogenation catalyst such as, for example, a ruthenium-BINAP complex, to produce the reduced product, substantially enriched in the more active isomer, represented by the formula:



wherein R', R³⁵ and R³⁶ represent radicals as defined above. Where the more active isomer has the R-stereochemistry, a Ru(R-BINAP) asymmetric hydrogenation catalyst can be utilized. Conversely, where the more active isomer has the S-stereochemistry, a Ru(S-BINAP) catalyst can be utilized. Where both isomers are active, or where it is desired to have a mixture of the two diastereomers, a hydrogenation catalyst such as platinum, or palladium, on carbon can be utilized to reduce the above compound. The reduced compound is then coupled to the urea isostere, as described above, to produce compounds of Formula III.

Preparation of Compounds of Formula IV

To produce compounds of Formula IV, starting with a lactate of the formula:




wherein P'' represents alkyl radicals, such as, for example, ethyl, methyl, benzyl and the like. The hydroxyl group of the lactate is protected as its ketal by reaction in a suitable solvent system with methyl isopropenyl ether (1,2-methoxypropene) in the presence of a suitable acid. Suitable solvent systems include methylene chloride, tetrahydrofuran and the like as well as mixtures thereof. Suitable acids include POCl₃ and the like. It should be noted that well-known groups other than methyl isopropenyl ether can be utilized to form the ketal. The ketal is then reduced with diisobutylaluminum hydride (DIBAL) at -78°C to produce the corresponding aldehyde which is then treated with ethylidene triphenylphosphorane (Wittig reaction) to produce a compound represented by the formula:

10

15


20




The ketal protecting group is then removed utilizing procedures well-known in the art such as by mild acid hydrolysis. The resulting compound is then esterified with isobutyryl chloride to produce a compound of the formula:

25


30



This compound is then treated with lithium diisopropyl amide at -78°C followed by warming of the reaction mixture to room temperature to effect a Claisen rearrangement ([3,3] sigmatropic rearrangement) to produce the corresponding acid represented by the formula:




Treatment of the acid with benzyl bromide in the presence of a tertiary amine base, e.g., DBU, produces the corresponding ester which is then cleaved oxidatively to give a trisubstituted succinic acid:



The trisubstituted succinic acid is then coupled to the urea isostere as described above. To produce the free acid, the benzyl ester is removed by hydrogenolysis to produce the corresponding acid. The acid can then be converted to the primary amide by methods well-known in the art.


An alternative method for preparing trisubstituted succinic acids involves reacting an ester of acetoacetic

acid represented by the formula:




5 where R is a suitable protecting group, such as methyl, ethyl, benzyl or t-butyl with sodium hydride and a hydrocarbyl halide ($R^{31}X$ or $R^{32}X$) in a suitable solvent, e.g., THF, to produce the corresponding disubstituted derivative represented by the formula:

10




This disubstituted acetoacetic acid derivative is then treated with lithium diisopropyl amide at about -10°C and in the presence of $\text{PhN}(\text{triflate})_2$ to produce a vinyl triflate of the formula:

15




20 The vinyl triflate is then carbonylated utilizing a palladium catalyst, e.g., $\text{Pd}_2(\text{OAc})_2(\text{Ph}_3\text{P})$, in the presence of an alcohol ($R''\text{OH}$) or water ($R''=\text{H}$) and a base, e.g., triethylamine, in a suitable solvent such as DMF, to produce the olefinic ester or acid of the formula:

25




The olefin can then be subsequently asymmetrically hydrogenated, as described above, to produce a trisubstituted succinic acid derivative of the formula:




If R'' is not H, R'' can be removed by either hydrolysis, acidolysis, or hydrogenolysis, to afford the corresponding acid, which is then coupled to the urea isostere as described above and then, optionally, the R group removed to produce the corresponding acid, and optionally, converted to the amide.

Alternatively, one can react the urea isostere with either a suitably monoprotected succinic acid or glutaric acid of the following structure;



followed by removal of the protecting group and conversion of the resulting acid to an amide. One can also react an anhydride of the following structure;



with the urea isostere and then separate any isomers or

convert the resulting acid to an amide and then separate any isomers.

It is contemplated that for preparing compounds of the Formulas having R^6 , the compounds can be prepared following the procedure set forth above and, prior to coupling the urea derivative or analog thereof, e.g. coupling to the amino acid $\text{PNH}(\text{CH}_2)_t\text{CH}(\text{R}^1)\text{COOH}$, carried through a procedure referred to in the art as reductive amination. Thus, a sodium cyanoborohydride and an appropriate aldehyde $\text{R}^6\text{C}(\text{O})\text{H}$ or ketone $\text{R}^6\text{C}(\text{O})\text{R}^6$ can be reacted with the urea derivative compound or appropriate analog at room temperature in order to reductively aminate any of the compounds of Formulas I-IV. It is also contemplated that where R^3 of the amino alcohol intermediate is hydrogen, the inhibitor compounds of the present invention wherein R^3 is alkyl, or other substituent wherein the $\alpha\text{-C}$ contains at least one hydrogen, can be prepared through reductive amination of the final product of the reaction between the amino alcohol and the amine or at any other stage of the synthesis for preparing the inhibitor compounds.

Contemplated equivalents of the general formulas set forth above for the antiviral compounds and derivatives as well as the intermediates are compounds otherwise corresponding thereto and having the same general properties such as tautomers thereof and compounds wherein one or more of the various R groups are simple variations of the substituents as defined therein, e.g., wherein R is a higher alkyl group than that indicated. In addition, where a substituent is designated as, or can be, a hydrogen, the exact chemical nature of a substituent which is other than hydrogen at that position, e.g., a hydrocarbyl radical or a halogen, hydroxy, amino and the like functional group, is not critical so long as it does not adversely affect the overall activity and/or synthesis procedure.

The chemical reactions described above are generally disclosed in terms of their broadest application to the preparation of the compounds of this invention.

5 Occasionally, the reactions may not be applicable as described to each compound included within the disclosed scope. The compounds for which this occurs will be readily recognized by those skilled in the art. In all such cases, either the reactions can be successfully performed by
10 conventional modifications known to those skilled in the art, e.g., by appropriate protection of interfering groups, by changing to alternative conventional reagents, by routine modification of reaction conditions, and the like, or other reactions disclosed herein or otherwise conventional, will
15 be applicable to the preparation of the corresponding compounds of this invention. In all preparative methods, all starting materials are known or readily preparable from known starting materials.

20 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of
25 the remainder of the disclosure in any way whatsoever.

All reagents were used as received without purification. All proton and carbon NMR spectra were obtained on either a Varian VXR-300 or VXR-400 nuclear
30 magnetic resonance spectrometer.

Example 1

Preparation of [1S-[1R*(R*), 2S*]]- N-[3-[[[(1,1-
dimethylethyl)aminolcarbonyl](2-methylpropyl)aminol -2-
5 hydroxy-1-(phenylmethyl)propyl]-2-[(2-
quinolinylcarbonyl)aminol]-butanediamide

Part A:

To a solution of 75.0g (0.226 mol) of N-
10 benzyloxycarbonyl-L-phenylalanine chloromethyl ketone in a
mixture of 807 mL of methanol and 807 mL of tetrahydrofuran
at -2°C, was added 13.17g (0.348 mol, 1.54 equiv.) of solid
sodium borohydride over one hundred minutes. The solvents
were removed under reduced pressure at 40°C and the residue
15 dissolved in ethyl acetate (approx. 1L). The solution was
washed sequentially with 1M potassium hydrogen sulfate,
saturated sodium bicarbonate and then saturated sodium
chloride solutions. After drying over anhydrous magnesium
sulfate and filtering, the solution was removed under
20 reduced pressure. To the resulting oil was added hexane
(approx. 1L) and the mixture warmed to 60°C with swirling.
After cooling to room temperature, the solids were
collected and washed with 2L of hexane. The resulting
solid was recrystallized from hot ethyl acetate and hexane
25 to afford 32.3g (43% yield) of N-benzyloxycarbonyl-3(S)-
amino-1-chloro-4-phenyl-2(S)-butanol, mp 150-151°C and
M+Li+ = 340.

Part B:

To a solution of 6.52g (0.116 mol, 1.2 equiv.) of potassium hydroxide in 968 mL of absolute ethanol at room temperature, was added 32.3g (0.097 mol) of N-CBZ-3(S)-amino-1-chloro-4-phenyl-2(S)-butanol. After stirring for fifteen minutes, the solvent was removed under reduced pressure and the solids dissolved in methylene chloride. After washing with water, drying over magnesium sulfate, filtering and stripping, one obtains 27.9g of a white solid. Recrystallization from hot ethyl acetate and hexane afforded 22.3g (77% yield) of N-benzyloxycarbonyl-3(S)-amino-1,2(S)-epoxy-4-phenylbutane, mp 102-103°C and MH⁺ 298.

Part C:

A solution of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane (1.00g, 3.36 mmol) and isobutylamine (4.90g, 67.2 mmol, 20 equiv.) in 10 mL of isopropyl alcohol was heated to reflux for 1.5 hours. The solution was cooled to room temperature, concentrated in vacuo and then poured into 100 mL of stirring hexane whereupon the product crystallized from solution. The product was isolated by filtration and air dried to give 1.18g, 95% of N=[3(S)-phenylmethylcarbamoyl]amino-2(R)-hydroxy-4-phenylbutyl]N-[(2-methylpropyl)]amine mp 108.0-109.5°C, MH⁺ m/z = 371.

Part D:

A solution of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[(2-methylpropyl)]amine in 10 mL of tetrahydrofuran was treated with ~~tert~~-butylisocyanate (267 mg, 2.70 mmol) at room temperature for 5 minutes. The solvent was removed in vacuo and replaced with ethyl acetate. The ethyl acetate solution was washed with 5% citric acid, water, and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo to give

1.19g, 97% of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)-amino]-2-hydroxy-4-phenyl]-1-[(2-methylpropyl)]amino-2-(1,1-dimethyl)amino]carbonyl]butane, MH^+ m/z - 470.

5 Part E:

A solution of (1.00g, 2.21 mmol) [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(2-methylpropyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane in 20 mL of methanol was
10 hydrogenated over 10% palladium-on-carbon for 4 hours to give [2(R), 3(S)]-N-[[3-amino]-2-hydroxy-4-phenyl]-1-[(2-methylpropyl)amino-1-(1,1-dimethylethyl)amino]carbonyl]butane 720 mg, 97%.

15 Part F:

A solution of N-Cbz-L-asparagine (602mg, 2.26 mmol) and N-hydroxybenzotriazole (493 mg, 3.22 mmol) in 2mL of dimethylformamide was cooled to 0°C and treated with EDC (473 mg, 2.47 mmol). The solution was allowed to stir at
20 0°C for 20 minutes and then treated with [2(R), 3(S)]-N-[[3-amino]-2-hydroxy-4-phenyl]-1-[(2-methylpropyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane (720 mg, 2.15 mmol) in 1mL of dimethylformamide. The solution was allowed to warm to room temperature and held at this
25 temperature for 7 hours. The reaction mixture was then poured into 100 mL of 60% saturated aqueous sodium bicarbonate whereupon a white precipitate formed that was isolated by filtration. The filter cake was washed with water, 5% aqueous citric acid, water and then dried in
30 vacuo to give 1.04g, 83% of [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]-carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide, mp: 164.0-166.5°C, MH^+ m/z = 584.

Part G.

A solution of [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)-amino]-butanediamide (1.00g, 1.72 mmol) in 10 mL of methanol was hydrogenated over 10% palladium-on-carbon for 4 hours to give [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-amino]-butanediamide, 784mg, 99%.

Part H:

A mixture of [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-amino]-butanediamide, (784 mg, 1.70 mmol), 2-quinoline carboxylic acid N-hydroxysuccinimide ester (459 mg, 1.70 mmol), N-methylmorpholine (343 mg, 3.40 mmol) in 5 mL of dichloromethane was stirred at room temperature for 15 minutes. The solvent was removed *in vacuo* and replaced with ethyl acetate and the solution washed with 5% aqueous citric acid, saturated aqueous sodium bicarbonate, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. The crude product was recrystallized from acetone/hexane to give 790 mg, 77% of [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide, mp 107.0-109.8°C, MH⁺ = 605.

Example 2

The procedure described in Example 1, part C-H, was used to prepare [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)-amino]-butanediamide.

a) From the reaction of 1.06g (3.56mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 6.25g (71.7mmol) of isoamylamine, one obtains 1.27g (92%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[(3-methylbutyl)]amine, mp 130-132°C and MH⁺ 385. This amine (400mg, 1.04mmol) was then reacted with tert-butylisocyanate (110mg, 1.11mmol) to afford 500mg (100%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(3-methylbutyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane; as an oil, MH⁺ 484.

b) The CBZ protected compound (530mg, 1.10mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (377mg, 1.42mmol) in the presence of N-hydroxybenzotriazole (290mg, 2.15mmol) and EDC (300mg, 1.56mmol) to yield 430mg (53%) of [1S-[1R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide, mp 148-151°C (dec) and MH⁺ 598. This compound (370mg, 0.619mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with 2-quinolinecarboxylic acid N-hydroxy-succinimide ester (193mg, 0.714mmol), in the presence of N-methylmorpholine, to afford 310mg (70%) of pure [1S-[1R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide; mp 93.5-95.5°C and MH⁺ 619.

Example 3

The procedure described in Example 1, part C-H, was used to prepare [1S-[1R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl]2-naphthylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amio]-butanediamide.

a) From the reaction of 1.80g (6.05mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 1.15g (7.31mmol) of 2-(aminomethyl)naphthalene, one obtains 2.11g (77%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[(2-naphthylmethyl)]amine, MH⁺ 455. This amine (366.8mg, 0.807mmol) was then reacted with tert-butylisocyanate (66.4mg, 0.67mmol) to afford 350.0mg (94%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(2-naphthylmethyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane; as an oil, MH⁺ 554.

b) The CBZ protected compound (330mg, 0.596mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (165.1mg, 0.62mmol) in the presence of N-hydroxybenzotriazole (142.3mg, 0.93mmol) and EDC (130.7mg, 0.68mmol) to yield 161.7mg (41%) of [1S-[1R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-naphthylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide; mp 151-152°C (dec) and MH⁺ 668. This compound (91.0mg, 0.136mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with 2-quinolinecarboxylic acid N-hydroxysuccinimide ester (36.8mg, 0.136mmol), in the presence of N-methylmorpholine, to afford

65.8mg (70%) of pure [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-naphthylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide; mp 119-120°C and
5 MH⁺ 689.

Example 4

The procedure described in Example 1, part C-H,
10 was used to prepare [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-phenylethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide.

15 a) From the reaction of 1.00g (3.36mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 8.19g (67.0mmol) of 2-phenethyl amine, one obtains 1.10g (79%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[(2-phenylethyl)]amine, mp 137-138°C
20 and MH⁺ 419. This amine (750mg, 1.79mmol) was then reacted with tert-butylisocyanate (178mg, 1.79mmol) to afford 897mg • (97%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(2-phenylethyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane; as an oil, MH⁺ 518.

25

b) The CBZ protected compound (897mg, 1.73mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (620.7mg, 2.33mmol) in the presence of N-
30 hydroxybenzotriazole (509.5mg, 3.33mmol) and EDC (488.0mg, 2.55mmol) to yield 1.00g (92%) of [1S-[1R*(R*), 2S*]]- N¹[3[[[(1,1-dimethylethyl)amino]carbonyl](2-phenylethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide; mp 145°C (dec)
35 and MH⁺ 632. This compound (860mg, 1.36mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with

2-quinolinecarboxylic acid N-hydroxysuccinimide ester (338mg, 1.25mmol), in the presence of N-methylmorpholine, to afford 450.4mg (55%) of pure [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-phenylethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide; mp 139-140°C and MH⁺ 653.

Example 5

The procedure described in Example 1, part C-H, was used to prepare [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2,2-dimethylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide.

a) From the reaction of 1.00g (3.36mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 7.9mL (approx. 67mmol) of neopentyl amine, one obtains 0.69g (49%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[(2,2-dimethylpropyl)]amine, MH⁺ 385. This amine (686mg, 1.78mmol) was then reacted with tert-butylisocyanate (180mg, 1.78mmol) to afford 860mg (100%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(2,2-dimethylpropyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane; MH⁺ 484.

b) The CBZ protected compound (860mg, 1.78mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (471mg, 1.77mmol) in the presence of N-hydroxybenzotriazole (406mg, 2.66mmol) and EDC (374mg, 1.95mmol) to yield 326mg (34%) of [1S-[R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2,2-dimethylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide; mp 177-178°C

and MH⁺ 598. This compound (245mg, 0.41mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with 2-quinolinecarboxylic acid N-hydroxy-succinimide ester (111mg, 0.41mmol), in the presence of N-methylmorpholine, to afford 150mg (59%) of pure [1S-[R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](2,2-dimethylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide; mp 115-117°C and MH⁺ 619.

Example 6

The procedure described in Example 1, part C-H, was used to prepare [1S-[R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-methoxyphenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide;

a) From the reaction of 1.00g (3.36mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 9.2g (67mmol) of 4-methoxybenzyl amine, one obtains 1.12g (76%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)-amino]-2-hydroxy-4-phenylbutyl]N-[(4-methoxyphenylmethyl)]amine, MH⁺ 435. This amine (1.12g, 2.58mmol) was then reacted with tert-butylisocyanate (260mg, 2.58mmol) to afford 1.35g (98%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(4-methoxyphenylmethyl)]amino-1-(1,1-dimethylethyl)amino]-carbonyl]butane; MH⁺ 534.

b) The CBZ protected compound (1.35g, 2.53mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (684mg, 2.57mmol) in the presence of N-hydroxybenzotriazole (590mg, 3.85mmol) and EDC (543mg,

2.83mmol) to yield 442mg (29%) of [1S-[1R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-methoxyphenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[phenylmethylcarbamoyl)amino]-butanediamide; mp 175°C (dec) and MH⁺ 648. This compound (345mg, 0.53mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with 2-quinolinecarboxylic acid N-hydroxy-succinimide ester (118mg, 0.44mmol), in the presence of N-methylmorpholine, to afford 108mg (31%) of pure [1S-[1R*(R*), 2S*]]-N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-methoxyphenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide; mp 220°C (dec) and MLi⁺ 675.

Example 7

The procedure described in Example 1, part C-H, was used to prepare [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](n-butyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide.

a) From the reaction of 1.48g (5.0mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 7.314g (100.0mmol) of n-butyl amine, one obtains 1.50g (80%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[n-butyl]amine. This amine (1.48g, 4.0mmol) was then reacted with tert-butylisocyanate (396mg, 4.0mmol) to afford 1.87g (100%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(n-butyl)amino-1-(1,1-dimethylethyl)amino]carbonyl] butane as an oil.

b) The CBZ protected compound (1.87g, 4.0mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (1.05g, 3.96mmol) in the presence of N-hydroxybenzotriazole (535mg, 7.9mmol) and EDC (759mg, 3.96mmol) to yield 1.75g (76%) of [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](n-butyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide; mp 166-167°C and MH⁺ 584.

Example 8

The procedure described in Example 1, part C-H, was used to prepare [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide.


a) From the reaction of 1.48g (5.0mmol) of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane and 10.68g (100.0mmol) of benzyl amine, one obtains 1.88g (95%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenylbutyl]N-[(phenylmethyl)]amine. This amine (1.88g, 4.65mmol) was then reacted with tert-butylisocyanate (460.0mg, 4.6mmol) to afford 2.24g (96%) of [2(R), 3(S)]-N-[[3-(phenylmethylcarbamoyl)amino]-2-hydroxy-4-phenyl]-1-[(phenylmethyl)]amino-1-(1,1-dimethylethyl)amino]carbonyl]butane.

b) The CBZ protected compound (2.22g, 4.4mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with N-CBZ-L-asparagine (1.17g, 4.4mmol) in the presence of N-hydroxybenzotriazole (1.19g, 8.8mmol) and EDC (843mg, 4.4mmol) to yield 2.11g (78%) of [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(phenylmethylcarbamoyl)amino]-butanediamide; mp 156-158°C and MH⁺ 618. This compound (1.0g, 1.62mmol) was then deprotected by hydrogenation over 10% palladium-on-carbon and the resulting free amine coupled with 2-quinolinecarboxylic acid N-hydroxysuccinimide ester (437mg, 1.62mmol), in the presence of N-methylmorpholine, to afford 640mg (62%) of pure [1S-[1R*(R*), 2S*]]- N¹[3-[[[(1,1-dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide; mp 110.5-112.5°C and MH⁺ 639.

Example 9

Additional exemplary compounds of the present invention are listed in Table 1. These compounds were prepared according to the following general procedures.
5 General Procedure for the Synthesis of 1,3-Diamino 4-phenyl Butan-2-ol Derivatives.

A mixture of the amine R^3NH_2 (20 equiv.) in dry isopropyl alcohol (20mL/mmol of epoxide to be converted) was heated to reflux and then treated with an N-Cbz amino epoxide of the formula:



15 from a solids addition funnel over a 10-15 minute period. After the addition is complete the solution was maintained at reflux for an additional 15 minutes and the progress of the reaction monitored by TLC. The reaction mixture was then concentrated in vacuo to give an oil that was treated with n-hexane with rapid stirring whereupon the ring opened material precipitated from solution. Precipitation was generally complete within 1 hr and the product was then isolated by filtration on a Büchner funnel and then air dried. The product was further dried in vacuo. This method
25 affords amino alcohols of sufficient purity for most purposes.

General Procedure for the Synthesis of Amino Epoxides

The amino epoxides can be prepared as shown in the previous example. Preferably, such amino epoxides are prepared
5 utilizing the following procedure.

Step A:

A solution of L-phenylalanine (50.0 g, 0.302 mol), sodium hydroxide (24.2 g, 0.605 mol) and potassium carbonate (83.6 g, 0.605 mol) in water (500 ml) is heated
10 to 97°C. Benzyl bromide (108.5 ml, 0.912 mol) is then slowly added (addition time ~25 min). The mixture is then stirred at 97°C for 30 minutes. The solution is cooled to room temperature and extracted with toluene (2 x 250 ml).
15 The combined organic layers are then washed with water, brine, dried over magnesium sulfate, filtered and concentrated to give an oil product. The crude product is then used in the next step without purification.

20 Step B:

The crude benzylated product of the above step is dissolved in toluene (750 ml) and cooled to -55°C. A 1.5 M solution of DIBAL-H in toluene (443.9 ml, 0.666 mol) is then added at a rate to maintain the temperature between
25 -55° to -50°C (addition time - 1 hour). The mixture is stirred for 20 minutes at -55°C. The reaction is quenched at -55°C by the slow addition of methanol (37 ml). The cold solution is then poured into cold (5°C) 1.5 N HCl solution (1.8 L). The precipitated solid (approx. 138 g)
30 is filtered off and washed with toluene. The solid material is suspended in a mixture of toluene (400 ml) and water (100 ml). The mixture is cooled to 5°C, treated with 2.5 N NaOH (186 ml) and then stirred at room temperature until the solid is dissolved. The toluene layer is
35 separated from the aqueous phase and washed with water and brine, dried over magnesium sulfate, filtered and concentrated to a volume of 75 ml (89 g). Ethyl acetate (25 ml) and hexane (25 ml) are then added to the residue

upon which the alcohol product begins to crystallize. After 30 min., an additional 50 ml hexane is added to promote further crystallization. The solid is filtered off and washed with 50 ml hexane to give approximately 35 g of material. A second crop of material can be isolated by refiltering the mother liquor. The solids are combined and recrystallized from ethyl acetate (20 ml) and hexane (30 ml) to give, in 2 crops, approximately 40 g (40% from L-phenylalanine) of analytically pure alcohol product. The mother liquors are combined and concentrated (34 g). The residue is treated with ethyl acetate and hexane which provides an additional 7 g (~7% yield) of slightly impure solid product. Further optimization in the recovery from the mother liquor is probable.

15

Step C:

A solution of oxalyl chloride (8.4 ml, 0.096 mol) in dichloromethane (240 ml) is cooled to -74°C. A solution of DMSO (12.0 ml, 0.155 mol) in dichloromethane (50 ml) is then slowly added at a rate to maintain the temperature at -74°C (addition time ~1.25 hr). The mixture is stirred for 5 min. followed by addition of a solution of the alcohol (0.074 mol) in 100 ml of dichloromethane (addition time ~20 min., temp. -75°C to -68°C). The solution is stirred at -78°C for 35 minutes. Triethylamine (41.2 ml, 0.295 mol) is then added over 10 min. (temp. -78°C to -68°C) upon which the ammonium salt precipitates. The cold mixture is stirred for 30 min. and then water (225 ml) is added. The dichloromethane layer is separated from the aqueous phase and washed with water, brine, dried over magnesium sulfate, filtered and concentrated. The residue is diluted with ethyl acetate and hexane and then filtered to further remove the ammonium salt. The filtrate is concentrated to give the desired aldehyde product. The aldehyde was carried on to the next step without purification.

Temperatures higher than -70°C has been reported in the literature for the Swern oxidation. Other Swern modifications and alternatives to the Swern oxidations are also possible.

5

A solution of the crude aldehyde 0.074 mol and chloriodomethane (7.0 ml, 0.096 mol) in tetrahydrofuran (285 ml) is cooled to -78°C . A 1.6 M solution of n-butyllithium in hexane (25 ml, 0.040 mol) is then added at a rate to maintain the temperature at -75°C (addition time - 15 min.). After the first addition, additional chloriodomethane (1.6 ml, 0.022 mol) is added again, followed by n-butyllithium (23 ml, 0.037 mol), keeping the temperature at -75°C . The mixture is stirred for 15 min. Each of the reagents, chloriodomethane (0.70 ml, 0.010 mol) and n-butyllithium (5 ml, 0.008 mol) are added 4 more times over 45 min. at -75°C . The cooling bath is then removed and the solution warmed to 22°C over 1.5 hr. The mixture is poured into 300 ml of saturated aq. ~~ammonium~~ chloride solution. The tetrahydrofuran layer is separated. The aqueous phase is extracted with ethyl acetate (1 x 300 ml). The combined organic layers are washed with brine, dried over magnesium sulfate, filtered and concentrated to give a brown oil (27.4 g). The product could be used in the next step without purification. The desired diastereomer can be purified by recrystallization at a subsequent step.

Alternately, the product could be purified by chromatography.

General procedure for the Reaction of Amino Alcohols with Isocyanates: Preparation of Ureas

A solution from the amino alcohol in tetrahydrofuran (THF) was treated at room temperature with the appropriate isocyanate of formula R^4NCO via syringe under nitrogen. After the reaction has stirred for ~5m the

progress of the reaction was monitored by TLC. The solvent was removed in vacuo and the product obtained was of sufficient purity for most purposes. The product may be further purified by dissolution in ethyl acetate and washing with 5% aqueous citric acid, water, and brine. The solvent is dried over anhydrous magnesium sulfate, filtered and concentrated in vacuo to give the pure urea.

General Procedure for the Removal of the Protecting Groups by Hydrogenolysis with Palladium on Carbon

A. Alcohol Solvent

The Cbz-protected peptide derivative was dissolved in methanol (ca.20mL/mmol) and 10% palladium on carbon catalyst is added under a nitrogen atmosphere. The reaction vessel is sealed and flushed 5 times with nitrogen and then 5 times with hydrogen. The pressure is maintained at 50 psig for 1-16 hours and then the hydrogen replaced with nitrogen and the solution filtered through a pad of celite to remove the catalyst. The solvent is removed in vacuo to give the free amino derivative of suitable purity to be taken directly on to the next step.

B. Acetic Acid Solvent

The Cbz-protected peptide derivative was dissolved in glacial acetic acid (20mL/mmol) and 10% palladium on carbon catalyst is added under a nitrogen atmosphere. The reaction vessel is flushed 5 times with nitrogen and 5 times with hydrogen and then maintained at 40 psig for about 2h. The hydrogen was then replaced with nitrogen and the reaction mixture filtered through a pad of celite to remove the catalyst. The filtrate was concentrated and the resulting product taken up in anhydrous ether and evaporated to dryness 3 times. The final product, the acetate salt, was dried in vacuo and is of suitable purity for subsequent conversion.

General Procedure for Removal of Boc-protecting Group with 4N Hydrochloric Acid in Dioxane


The Boc-protected amino acid or peptide is treated with a solution of 4N HCl in dioxane with stirring at room temperature. Generally the deprotection reaction is complete within 15 minutes, the progress of the reaction is monitored by thin layer chromatography (TLC). Upon completion, the excess dioxane and HCl are removed by evaporation in vacuo. The last traces of dioxane and HCl are best removed by evaporation again from anhydrous ether or acetone. The hydrochloride salt thus obtained is thoroughly dried in vacuo and is suitable for further reaction.

15 EDC/HOBt Coupling of Cbz-Asparagine (General Procedure)

N-CBZ-(L-asparagine (1.10eq) and N-hydroxybenzotriazole (HOBt) (1.5eq) are dissolved in dry dimethylformamide (DMF) (2-5mL/mmol) and cooled in an ice bath. 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (1.10eq) is added to the stirring solution and maintained at 0°C for 10 minutes. A solution of the amino component (free amine), 1.0eq in DMF (1-2mL/mmol), is added. [In the case of the amine hydrochloride or acetate salt, an equivalent of N-methylmorpholine is also added.] The reaction mixture is stirred at 0°C for 1 hour and then at room temperature for ~5-6 hours. The reaction mixture is then poured into a rapidly stirring solution of 60% saturated aqueous sodium bicarbonate (ca-50mL/mmol). An immediate white precipitate forms which is collected on a Büchner funnel and the solid washed thoroughly with saturated aqueous sodium bicarbonate, water, 5% aqueous citric acid solution and water. The product is thoroughly dried in vacuo and redissolved in DMF, filtered and reprecipitated by the addition to water. The precipitated product is isolated by filtration, washed again with water and dried in vacuo.

General Procedure for Acylation with 2-Quinoline Carboxylic
Acid N-Hydroxysuccinimide Ester

5 A solution of the free amine (or amine acetate
salt) and 1.0 equivalent of N-hydroxysuccinimide 2-quinoline
carboxylate in anhydrous dichloromethane was treated with
1.5 equivalents of N-methylmorpholine (NMM) at room
temperature. The progress of the reaction was monitored by
10 TLC and when the reaction was complete the reaction mixture
was diluted with additional dichloromethane and the solution
washed with saturated aqueous sodium bicarbonate, 5% aqueous
citric acid, water and brine. The solution was dried over
anhydrous magnesium sulfate, filtered and concentrated in
15 vacuo. The product thus obtained was recrystallized from a
mixture of acetone and hexane.

TABLE 1

5

Entry No.	R	R ³	R ⁴
10	1 Cbz ^a	CH ₃	n-Butyl
	2 Cbz	i-Butyl	CH ₃
	3 Cbz	i-Butyl	n-Butyl
	4 Q ^b	i-Butyl	n-Butyl
	5 Cbz	i-Propyl	n-Butyl
15	6 Q	i-Propyl	n-Butyl
	7 Cbz	C ₆ H ₅	n-Butyl
	8 Cbz	-CH ₂ -	n-Butyl
	9 Cbz	-CH ₂ -	n-Butyl
	10 Q	-CH ₂ -	n-Butyl
20	11 Cbz		n-Butyl
	12 Cbz	i-Butyl	n-Propyl
	13 Cbz	i-Butyl	-CH ₂ CH(CH ₃) ₂
	14 Cbz	(R)-CH(CH ₃)-	n-Butyl
	15 Cbz	-CH ₂ -	i-Propyl

TABLE 1 (Cont'd.)

Entry No.	R	R ³	R ⁴	
5	16	Cbz	$-\text{CH}_2-\text{C}_6\text{H}_{11}$	$-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$
	17	Cbz	i-Butyl	$-\text{CH}_2\text{CH}_3$
	18	Cbz	i-Butyl	$-\text{CH}(\text{CH}_3)_2$
	19	Cbz	i-Butyl	$-\text{C}_6\text{H}_{11}$
	20	Q	i-Butyl	$-\text{C}_6\text{H}_{11}$
10	21	Cbz	$-\text{CH}_2-\text{C}_6\text{H}_{11}$	$-(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2$
	22	Cbz	$(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2$	$-\text{CH}(\text{CH}_3)_2$
	23	Q	i-Butyl	$-\text{CH}(\text{CH}_3)_2$
	24	Cbz	i-Butyl	$-\text{C}(\text{CH}_3)_3$
	25	Q	i-Butyl	$-\text{C}(\text{CH}_3)_3$
15	26	Cbz	$-\text{CH}_2-\text{C}_{10}\text{H}_7$	$-\text{C}(\text{CH}_3)_3$
	27	Q	$-\text{CH}_2-\text{C}_{10}\text{H}_7$	$-\text{C}(\text{CH}_3)_3$
	28	Cbz	$-(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2$	$-\text{C}(\text{CH}_3)_3$
	29	Q	$-(\text{CH}_2)_2\text{CH}(\text{CH}_3)_2$	$-\text{C}(\text{CH}_3)_3$
	30	Cbz	$-\text{CH}_2\text{C}_6\text{H}_5$	$-\text{C}(\text{CH}_3)_3$
20	31	Q	$-\text{CH}_2\text{C}_6\text{H}_5$	$-\text{C}(\text{CH}_3)_3$
	32	Cbz	$-(\text{CH}_2)_2\text{C}_6\text{H}_5$	$-\text{C}(\text{CH}_3)_3$
	33	Cbz	$-(\text{CH}_2)_2\text{C}_6\text{H}_5$	$-\text{C}(\text{CH}_3)_3$
	34	Cbz	n-Butyl	$-\text{C}(\text{CH}_3)_3$
	35	Cbz	n-Pentyl	$-\text{C}(\text{CH}_3)_3$
25	36	Cbz	n-Hexyl	$-\text{C}(\text{CH}_3)_3$
	37	Cbz	$-\text{CH}_2-\text{C}_6\text{H}_5$	$-\text{C}(\text{CH}_3)_3$
	38	Cbz	$-\text{CH}_2\text{C}(\text{CH}_3)_3$	$-\text{C}(\text{CH}_3)_3$
	39	Q	$-\text{CH}_2\text{C}(\text{CH}_3)_3$	$-\text{C}(\text{CH}_3)_3$
	40	Cbz	$-\text{CH}_2\text{CH}_2-\text{N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{O}$	$-\text{C}(\text{CH}_3)_3$
30	41	Cbz	$-\text{CH}_2\text{C}_6\text{H}_5\text{OCH}_3$ (para)	$-\text{C}(\text{CH}_3)_3$

TABLE 1 (Cont'd.)











Entry No.	R	R ³	R ⁴
5 42	Cbz		-C(CH ₃) ₃
43	Cbz		-C(CH ₃) ₃
44	Cbz	-(CH ₂) ₂ C(CH ₃) ₃	-C(CH ₃) ₃
45	Q	-(CH ₂) ₂ C(CH ₃) ₃	-C(CH ₃) ₃
46	Cbz	-(CH ₂) ₄ OH	-C(CH ₃) ₃
10 47	Q	-(CH ₂) ₄ OH	-C(CH ₃) ₃
48	Q		-C(CH ₃) ₃
49	Q		-C(CH ₃) ₃
50		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
51		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
15 52		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
53		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
54		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
55		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃

TABLE 1 (Cont'd.)







Entry No.	R	R ³	R ⁴
5 56		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
57		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
58		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
59		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
60		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
10 61		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃

TABLE 1 (Cont'd.)









Entry No.	R	R ³	R ⁴
5 62		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
63		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
64		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
65		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
66		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
10 67		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃


TABLE 1 (Cont'd.)

Entry No.	R	R ³	R ⁴
5	<hr/>		
68		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
69		-CH ₂ CH(CH ₃) ₂	-C(CH ₃) ₃
10	<hr/>		
	a benzyloxycarbonyl		
	b 2-quinolinylcarbonyl		

EXAMPLE 10

Following the generalized procedures set forth in Example 9, the compounds set forth in Table 2 were prepared.

TABLE 2



10



Entry	A	R ³	R ⁴	
15	1	Cbz-Val	<i>i</i> -amyl	tBu
	2	Cbz-Leu	<i>i</i> -amyl	<i>t</i> -Bu
	3	Cbz-Ile	<i>i</i> -amyl	<i>t</i> -Bu
	4	Ac-D- <u>homo</u> -Phe	<i>i</i> -Bu	<i>n</i> -Bu
20	5	Qui-Orn (γ-Cbz)	-CH ₂ - 	<i>t</i> -Bu
	6	Cbz-Asn	-CH ₂ CH=CH ₂	<i>t</i> -Bu
	7	Acetyl- <i>t</i> -BuGly	<i>i</i> -amyl	<i>t</i> -Bu
	8	Acetyl-Phe	<i>i</i> -amyl	<i>t</i> -Bu
	9	Acetyl-Ile	<i>i</i> -amyl	<i>t</i> -Bu
25	10	Acetyl-Leu	<i>i</i> -amyl	<i>t</i> -Bu
	11	Acetyl-His	<i>i</i> -amyl	<i>t</i> -Bu
	12	Acetyl-Thr	<i>i</i> -amyl	<i>t</i> -Bu
	13	Acetyl-NHCH(C(CH ₃) ₂ (SCH ₃))C(O)-	<i>i</i> -amyl	<i>t</i> -Bu
30	14	Cbz-Asn	<i>i</i> -amyl	<i>t</i> -Bu
	15	Cbz-Ala	<i>i</i> -amyl	<i>t</i> -Bu

TABLE 2 (Cont'd)

	Entry	A	R ³	R ⁴
5				
	16	Cbz-Ala	<u>i</u> -amyl	<u>t</u> -Bu
	17	Cbz-beta-cyanoAla	<u>i</u> -amyl	<u>t</u> -Bu
	18	Cbz-t-BuGly	<u>i</u> -amyl	<u>t</u> -Bu
10	19	Q-t-BuGly	<u>i</u> -amyl	<u>t</u> -Bu
	20	Q-SCH ₃ Cys	<u>i</u> -amyl	<u>t</u> -Bu
	21	Cbz-SCH ₃ Cys	<u>i</u> -amyl	<u>t</u> -Bu
	22	Q-Asp	<u>i</u> -amyl	<u>t</u> -Bu
	23	Cbz-(NHCH(C(CH ₃) ₂ (SCH ₃))C(O)	<u>i</u> -amyl	<u>t</u> -Bu
15	24	Cbz-EtGly	<u>i</u> -amyl	<u>t</u> -Bu
	25	Cbz-PrGly	<u>i</u> -amyl	<u>t</u> -Bu
	26	Cbz-Thr	<u>i</u> -amyl	<u>t</u> -Bu
	27	Q-Phe	<u>i</u> -amyl	<u>t</u> -Bu
	28	Cbz-Phe	<u>i</u> -amyl	<u>t</u> -Bu
20	29	Cbz [NHCH(CH ₂ C(O)NHCH ₃)C(O)	<u>i</u> -amyl	<u>t</u> -Bu
	30	Q[NHCH(CH ₂ C(O)NHCH ₃)C(O)	<u>i</u> -amyl	<u>t</u> -Bu

EXAMPLE 11

Following the generalized procedure of Example 9,
the compounds listed in Table 3 were prepared.

TABLE 3

10


Entry	R ¹
15	1 -CH ₂ SO ₂ CH ₃
	2 (R) -CH(OH)CH ₃
	3 -CH(CH ₃) ₂
	4 - (R, S) CH ₂ SOCH ₃
20	5 -CH ₂ SO ₂ NH ₂
	6 -CH ₂ SCH ₃
	7 -CH ₂ CH(CH ₃) ₂
	8 -CH ₂ CH ₂ C(O)NH ₂
	9 (-S) -CH(OH)CH ₃
25	10 -CH ₂ C≡CH
	11 -CF ₃
	12 -CH(CF ₃) ₂
	13 -C(CH ₃) ₂ (SCH ₃)

EXAMPLE 12

Following the generalized procedures of Example 9,
the compounds set forth in Table 4 were prepared.

5

TABLE 4



10

	Entry	R ²	A
15	1	n-Bu	Cbz-Asn
	2	cyclohexylmethyl	Cbz-Asn
	3	n-Bu	Boc
	4	n-Bu	Cbz
	5	C ₆ H ₅ CH ₂	Boc
20	6	C ₆ H ₅ CH ₂	Cbz
	7	C ₆ H ₅ CH ₂	benzoyl
	8	cyclohexylmethyl	Cbz
	9	n-Bu	Q-Asn
25	10	cyclohexylmethyl	Q-Asn
	11	C ₆ H ₅ CH ₂	Cbz-Ile
	12	C ₆ H ₅ CH ₂	Q-Ile
	13	C ₆ H ₅ CH ₂	Cbz-t-BuGly
	14	C ₆ H ₅ CH ₂	Q-t-BuGly
	15	C ₆ H ₅ CH ₂	Cbz-Val
30	16	C ₆ H ₅ CH ₂	Q-Val


TABLE 4 (Cont'd.)

Entry	R ²	A
5		
17	2-naphthylmethyl	Cbz-Asn
18	2-naphthylmethyl	Q-Asn
19	2-naphthylmethyl	Cbz
20	n-Bu	Cbz-Val
10	21	Q-Val
22	n-Bu	Q-Ile
23	n-Bu	Cbz-t-BuGly
24	n-Bu	Q-t-BuGly
25	p-F(C ₆ H ₄)CH ₂	Q-Asn
15	26	Cbz
27	p-F(C ₆ H ₄)CH ₂	Cbz-Asn
28	C ₆ H ₅ CH ₂	Cbz-propargylglycine
29	C ₆ H ₅ CH ₂	Q-propargylglycine
30	C ₆ H ₅ CH ₂	acetyl-propargylglycine
20	31	Cbz-Asn
	i-Bu	

EXAMPLE 13

The compounds listed in Table 5 were prepared according to the generalized procedures of Example 9.

5

TABLE 5

10


Entry	X(H)R ⁴	A
15	1	-NH ^t Bu
	2	-NEt ₂
	3	-NHC(CH ₃) ₂ CH ₂ CH ₃
		Cbz-Asn
		Cbz
		Cbz

EXAMPLE 14

The compounds of Table 6 were prepared according to the generalized procedures set forth in Example 9 except that instead of an isocyanate, an isothiocyanate equivalent was utilized.

TABLE 6

10



15

Entry

XHR⁴

1

NH₂Et

2

NH^tBu

20

The Cbz group of the compounds shown in Examples 13 and 14 can be removed as described in Example 9 and the resulting compound can be coupled to a desired α - or β -amino acid or the like to produce compounds of the present invention.

25


Example 15

The compounds shown in Table 7 were prepared according to the following general procedure.

- 5 This general procedure represents a Curtius Rearrangement and reaction with the amino alcohol derivative as prepared following the general procedure in Example 9.

To a solution of 1 mmol of carboxylic acid in 12
10 mL of toluene and 3 mmol of triethylamine at 90°C under a nitrogen atmosphere, was added 1 mmol of diphenylphosphoryl azide. After 1 hour, a solution of 1 mmol of amino alcohol derivative in 3.5 mL of either N,N-dimethylformamide or toluene was added. After 1 hour, the solvent was removed
15 under reduced pressure, ethyl acetate and water added and the layers separated. The organic layer was washed with 5% citric acid, sodium bicarbonate, brine, dried, filtered and concentrated to afford the crude product. This was then recrystallized or chromatographed on silica gel to afford the
20 purified final compound.

TABLE 7







5


R3

R4


10

$$-\text{CH}_2\text{CH}(\text{CH}_3)_2$$
$$-\text{C}(\text{CH}_3)_2$$
$$-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$$

$$-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$$

$$-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$$

$$-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$$


15

Example 16A. Preparation of 4(4-methoxybenzyl)itaconate


A 5 L three-necked round bottomed flask equipped with constant pressure addition funnel, reflux condenser, nitrogen inlet, and mechanical stirrer was charged with itaconic anhydride (660.8g, 5.88 mol) and toluene (2300 mL). The solution was warmed to reflux and treated with 4-methoxybenzyl alcohol (812.4g, 5.88 mol) dropwise over a 2.6h period. The solution was maintained at reflux for an additional 1.5h and then the contents were poured into three 2 L erlenmeyer flasks to crystallize. The solution was allowed to cool to room temperature whereupon the desired mono-ester crystallized. The product was isolated by filtration on a Buchner funnel and air dried to give 850.2g, 58% of material with mp 83-85°C, a second crop, 17% was isolated after cooling of the filtrate in an ice bath. ¹H NMR (CDCl₃) 300 MHz 7.32(d, J=8.7 Hz, 2H), 6.91(d, J=8.7 Hz, 2H), 6.49(s, 1H), 5.85(s, 1H), 5.12(s, 2H), 3.83(s, 3H), 3.40(s, 2H).

B. Preparation of Methyl 4(4-methoxybenzyl) itaconate

5 A 5 L three-necked round bottomed flask equipped with reflux condenser, nitrogen inlet, constant pressure addition funnel and mechanical stirrer was charged with 4(4-methoxybenzyl) itaconate (453.4g, 1.81 mol) and treated with 1,5-diazabicyclo[4.3.0]non-5-ene (275.6g, 1.81 mol), (DBN),
10 dropwise so that the temperature did not rise above 15°C. To this stirring mixture was added a solution of methyl iodide (256.9g, 1.81 mol) in 250 mL of toluene from the dropping funnel over a 45m period. The solution was allowed to warm to room temperature and stirred for an additional
15 3.25h.

The precipitated DBN hydroiodide was removed by filtration, washed with toluene and the filtrate poured into a separatory funnel. The solution was washed with sat. aq.
20 NaHCO_3 (2 X 500 mL), 0.2N HCl (1 X 500 mL), and brine (2 X 500 mL), dried over anhyd. MgSO_4 , filtered, and the solvent removed in vacuo. This gave a clear colorless oil, 450.2g, 94% whose NMR was consistent with the assigned structure.
 ^1H NMR (CDCl_3) 300 MHz 7.30(d, $J=8.7$ Hz, 2H), 6.90(d, $J=8.7$
25 Hz, 2H), 6.34(s, 1H), 5.71(s, 1H), 5.09(s, 2H), 3.82(s, 3H), 3.73(s, 3H), 3.38(s, 2H). ^{13}C NMR (CDCl_3) 170.46, 166.47, 159.51, 133.55, 129.97, 128.45, 127.72, 113.77, 66.36, 55.12, 51.94, 37.64.


C. Preparation of Methyl 4(4-methoxybenzyl) 2(R)-
methylsuccinate




5 A 500 mL Fisher-Porter bottle was charged with methyl 4(4-methoxybenzyl) itaconate (71.1g, 0.269 mol), rhodium (R,R) DiPAMP catalyst (204mg, 0.269 mmol, 0.1 mol%) and degassed methanol (215 mL). The bottle was flushed 5
10 times with nitrogen and 5 times with hydrogen to a final pressure of 40 psig. The hydrogenation commenced immediately and after ca. 1h the uptake began to taper off, after 3h the hydrogen uptake ceased and the bottle was flushed with nitrogen, opened and the contents concentrated
15 on a rotary evaporator to give a brown oil that was taken up in boiling *iso*-octane (ca. 200 mL, this was repeated twice), filtered through a pad of celite and the filtrate concentrated *in vacuo* to give 66.6g, 93% of a clear colorless oil, ¹H NMR (CDCl₃ 300 MHz 7.30(d, J=8.7 Hz, 2H),
20 6.91(d, J=8.7 Hz, 2H), 5.08(s, 2H), 3.82(s, 3H), 3.67(s, 3H), 2.95(ddq, J=5.7, 7.5, 8.7 Hz, 1H), 2.79(dd, J=8.1, 16.5 Hz, 1H), 2.45(dd, J=5.7, 16.5 Hz, 1H), 1.23(d, J=7.5 Hz, 3H).

D. Preparation of Methyl 2(R)-methylsuccinate

A 3 L three-necked round-bottomed flask equipped with a nitrogen inlet, mechanical stirrer, reflux condenser and constant pressure addition funnel was charged with methyl 4(4-methoxybenzyl) 2(R)-methylsuccinate (432.6g, 1.65 mol) and toluene (1200 mL). The stirrer was started and the solution treated with trifluoroacetic acid (600 mL) from the dropping funnel over 0.25h. The solution turned a deep purple color and the internal temperature rose to 45°C. After stirring for 2.25h the temperature was 27°C and the solution had acquired a pink color. The solution was concentrated on a rotary evaporator. The residue was diluted with water (2200 mL) and sat. aq. NaHCO₃ (1000 mL). Additional NaHCO₃ was added until the acid had been neutralized. The aqueous phase was extracted with ethyl acetate (2 X 1000 mL) to remove the by-products and the aqueous layer was acidified to pH=1.8 with conc. HCl. This solution was extracted with ethyl acetate (4 X 1000 mL), washed with brine, dried over anhyd. MgSO₄, filtered and concentrated on a rotary evaporator to give a colorless liquid 251g, >100% that was vacuum distilled through a short path apparatus cut 1: bath temperature 120°C @ >1mm, bp 25-29°C; cut 2: bath temperature 140°C @ 0.5mm, bp 95-108°C, 151g, [d]_D @ 25°C=+1.38°C (c=15.475, MeOH), [α]_D=+8.48°C (neat); cut 3: bath temperature 140°C, bp 108°C, 36g, [α]_D @ 25°C=+1.49°C (c=15.00, MeOH), [α]_D=+8.98°C (neat). Cuts 2 and 3 were combined to give 189g, 78% of product, ¹H NMR (CDCl₃) 300 MHz 11.6(brs, 1H), 3.72(s, 3H), 2.92(ddq, J=5.7, 6.9, 8.0 Hz, 1H), 2.81(dd, J=8.0, 16.8 Hz, 1H), 2.47(dd, J=5.7, 16.8 Hz, 1H), 1.26(d, J=6.9 Hz, 3H).

E. Preparation of Methyl Itaconate

5 A 50 mL round bottomed flask equipped with reflux condenser, nitrogen inlet and magnetic stir bar was charged with methyl 4(4-methoxybenzyl) itaconate (4.00g, 16 mmol). The solution was kept at room temperature for 18 hours and then the volatiles were removed in vacuo. The residue was
10 taken up in ethyl acetate and extracted three times with saturated aqueous sodium bicarbonate solution. The combined aqueous extract was acidified to pH=1 with aqueous potassium bisulfate and then extracted three times with ethyl acetate. The combined ethyl acetate solution was washed with
15 saturated aqueous sodium chloride, dried over anhydrous magnesium sulfate, filtered, and concentrated in vacuo. The residue was then vacuum distilled to give 1.23g, 75% of pure product, bp 85-87 @ 0.1 mm. ¹H NMR (CDCl₃) 300 MHz 6.34(s, 1H), 5.73(s, 2H), 3.76(s, 3H), 3.38(s, 2H). ¹³C NMR (CDCl₃)
20 177.03, 166.65, 129.220, 132.99, 52.27, 37.46.

F. Curtius Rearrangement of Methyl 2(R)-methylsuccinate:
Preparation of Methyl N-Moz-α-methyl β-alanine.


25

A 5L four necked round bottomed flask equipped with a nitrogen inlet, reflux condenser, mechanical stirrer, constant pressure addition funnel, and thermometer adapter
30 was charged with methyl 2(R)-methylsuccinate (184.1g, 1.26 mol), triethylamine (165.6g, 218 mL, 1.64 mol, 1.3 equivalents), and toluene (1063 mL). The solution was

warmed to 85°C and then treated dropwise with a solution of diphenylphosphoryl azide (346.8g, 1.26 mol) over a period of 1.2h. The solution was maintained at that temperature for an additional 1.0h and then the mixture was treated with 4-methoxybenzyl alcohol (174.1g, 1.26 mol) over a 0.33h period from the dropping funnel. The solution was stirred at 88°C for an additional 2.25h and then cooled to room temperature. The contents of the flask were poured into a separatory funnel and washed with sat. aq. NaHCO₃ (2 X 500 mL), 0.2N HCl (2 X 500 mL), brine (1 X 500 mL), dried over anhyd. MgSO₄, filtered, and concentrated in vacuo to give 302.3g, 85% of the desired product as a slightly brown oil. ¹H NMR (CDCl₃) 300 MHz 7.32(d, J=8.4 Hz, 2H), 6.91(d, J=8.4 Hz, 2H), 5.2(brm, 1H), 5.05(s, 2H), 3.83(s, 3H), 3.70(s, 3H), 3.35(m, 2H), 2.70(m, 2H), 1.20(d, J=7.2 Hz, 3H).

G. Hydrolysis of Methyl N-Moz-α-methyl β-alanine:

Preparation of α-methyl β-alanine Hydrochloride




A 5 L three-necked round bottomed flask equipped with a reflux condenser, nitrogen inlet and mechanical stirrer was charged with methyl N-Moz-α-methyl β-alanine (218.6g, 0.78 mol), glacial acetic acid (975 mL) and 12N hydrochloric acid (1960 mL). The solution was then heated to reflux for 3h. After the solution had cooled to room temperature (ca. 1h) the aqueous phase was decanted from organic residue (polymer) and the aqueous phase concentrated on a rotary evaporator. Upon addition of acetone to the concentrated residue a slightly yellow solid formed that was slurried with acetone and the white solid was isolated by filtration on a Buchner funnel. The last traces of acetone were removed by evacuation to give 97.7g, 90% of pure product, mp 128.5-130.5°C [α]_D @ 25°C=9.0°C (c=2.535,

Methanol). ^1H NMR (D_2O) 300 MHz 3.29(dd, $J=8.6$, 13.0 Hz, 1H), 3.16(dd, $J=5.0$, 13.0m Hz, 1H), 2.94(ddq, $J=7.2$, 5.0, 8.6 Hz, 1H), 1.30(d, $J=7.2$ Hz, 3H); ^{13}C NMR (D_2O) 180.84, 44.56, 40.27, 17.49.

5

H. Preparation of N-Boc α -Methyl β -Alanine



10 A solution of α -methyl β -alanine hydrochloride (97.7g, 0.70 mol) in water (1050 mL) and dioxane (1050 mL) the pH was adjusted to 8.9 with 2.9N NaOH solution. This stirring solution was then treated with di-tert- butyl pyrocarbonate (183.3g, 0.84 mol, 1.2 equivalents) all at
15 once. The pH of the solution was maintained between 8.7 and 9.0 by the periodic addition of 2.5N NaOH solution. After 2.5h the pH had stabilized and the reaction was judged to be complete. The solution was concentrated on a rotary evaporator (the temperature was maintained at $<40^\circ\text{C}$). The
20 excess di-tert -butyl pyrocarbonate was removed by extraction with dichloromethane and then the aqueous solution was acidified with cold 1N HCl and immediately extracted with ethyl acetate (4 X 1000 mL). The combined ethyl acetate extract was washed with brine, dried over anhyd. MgSO_4 ,
25 filtered and concentrated on a rotary evaporator to give a thick oil 127.3g, 90% crude yield that was stirred with n-hexane whereupon crystals of pure product formed, 95.65g, 67%, mp $76-78^\circ\text{C}$, $[\alpha]_D$ @ $25^\circ\text{C} = -11.8^\circ\text{C}$ ($c=2.4$, EtOH). A second crop was obtained by concentration of the filtrate and
30 dilution with hexane, 15.4g, for a combined yield of 111.05g, 78%. ^1H NMR (acetone D_6) 300 MHz 11.7 (brs, 1H), 6.05 (brs 1H), 3.35 (m, 1H), 3.22 (m, 1H), 2.50 (m, 1H), 1.45(s, 9H), 1.19 (d, $J=7.3$ Hz, 3H); ^{13}C NMR (acetone D_6) 177.01, 79.28, 44.44, 40.92, 29.08, 15.50. Elemental analysis calc'd. for

C₉H₁₇NO₄: C, 53.19, H, 8.42; N, 6.89. Found: C, 53.36; H, 8.46; N, 6.99.

5 I. Preparation of N-4-Methoxybenzyloxycarbonyl α -Methyl β -
5 Alanine

A solution of N-4-methoxybenzyloxycarbonyl
 α -methyl β -alanine methyl ester (2.81g, 10.0 mmol) in 30 mL
of 25% aqueous methanol was treated with lithium hydroxide
10 (1.3 equivalents) at room temperature for a period of 2h.
The solution was concentrated in vacuo and the residue taken
up in a mixture of water and ether and the phases separated
and the organic phase discarded. The aqueous phase was
acidified with aqueous potassium hydrogen sulfate to pH=1.5
15 and then extracted three times with ether. The combined
ethereal phase was washed with saturated aqueous sodium
chloride solution, dried over anhydrous magnesium sulfate,
filtered and concentrated in vacuo to give 2.60 g, 97% of N-
4-Methoxybenzyloxycarbonyl α -methyl β -alanine (N-Moz-AMBA)
20 which was purified by recrystallization from a mixture of
ethyl acetate and hexane to give 2.44g, 91% of pure product,
mp 96-97°C, MH+=268. ¹H NMR (D₆-acetone/300 MHz) 1.16 (3H, d,
J=7.2Hz), 2.70 (1H, m), 3.31 (2H, m), 3.31 (3H, s), 4.99 (2H,
s), 6.92 (2H, 4, J=8.7 Hz), 7.13 (2H, d, J=8.7 Hz).

25 J. Preparation of Propanamide, 3-(4-methoxybenzyl-
oxycarbonyl)-N-[3-[[[(1,1-dimethylethyl)aminyl]carbonyl](3-
methylbutyl)aminol-2-hydroxy-1-(phenylmethyl)propyl]-2-
methyl-[1S-[1R*(S*), 2S*]]-

30 . N-Moz-AMBA (468mg, 1.75mmol) was dissolved in 5mL
of DMF, HOBT (355mg, 2.6mmol) was added and the solution was
cooled to 0°C. The solution was treated with (336mg,
1.75mmol) EDC for 15 minutes. To this was added (612mg,
35 1.75mmol) of [2R,3S 3-amino-1-isoamyl-1-(t-
butylcarbonyl)amino 4-phenyl-2-butanol in 10mL of DMF and the
reaction stirred for 16 hours at room temperature. The DMF
was concentrated to 5mL and the product was precipitated by

addition to 60% saturated aqueous NaHCO₃. The solid was taken up in ethyl acetate and washed with KHSO₄, NaHCO₃, NaCl(saturated), dried over MgSO₄ and concentrated to yield 680mg of crude product which was crystallized from CH₂Cl₂, Et₂O, hexane, to yield 300mg of pure product.

Example 17

The compounds of Table 8 were prepared according to the procedure listed below and that utilized in Example 16.

Propaneamide, 3-[(1,1-dimethylethyl)butoxycarbonyl]amino-N-[3-[[[(1,1--dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]- (Table 8, Entry 11)

Part A:

A solution of N-t-butyloxycarbonyl-2-(R)-methyl-3-aminopropionic acid (372 mg, 1.83 mmol) and N-hydroxybenzotriazole (371 mg, 2.75 mmol) in 5 mL of dimethylformamide was cooled to 0°C. To this was added EDC (351 mg, 1.83 mmol) and the solution was stirred for 15 minutes. To this chilled solution was added a solution of 3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)-amino]-2(R)-hydroxy-1(S)(phenylmethyl)propylamine in 5 mL of dimethylformamide and stirred for 15 hours. The dimethylformamide was removed and replaced with 50 mL of ethyl acetate, and the organic phase was extracted with 5% potassium hydrogen sulfate, saturated sodium bicarbonate and brine. The ethyl acetate layer was dried over magnesium sulfate, filtered and concentrated to yield 613 mg of product after recrystallization from ethyl acetate, hexanes. (63 % yield). M+Li 541

Part B:

Preparation of Propaneamide, 3-amino-N-[3-[[[(1,1-dimethylethyl)amino] carbonyl]- (3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*), 2S*]-hydrochloride


The product from part A. (577 mg, 1.08 mmol) was dissolved in 40 mL of 4N HCl in dioxane and the solution stirred for 2 hours, and concentrated to yield the hydrochloride salt in quantitative yield.

Part C:

Preparation of Propaneamide, 3-(2-methylpropanoylamino)-N-[3-[[[(1,1-dimethylethyl)-amino]carbonyl] (3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*), 2S*]-

The product from part B. (236 mg, 0.5 mmol) was dissolved in anhydrous tetrahydrofuran and to this was added N-methylmorpholine (160 mg, 1.5 mmol) upon which time a precipitate formed. To this suspension was added isobutyryl chloride (53.5 mg, 0.5 mmol) and the suspension stirred for 15 hours. The suspension was diluted with ethyl acetate and washed with 5% potassium hydrogen sulfate, saturated sodium bicarbonate and brine. the organic layer was dried over magnesium sulfate, filtered and concentrated to yield 195 mg of crude product which was chromatographed on silica gel with 5% methanol methylene chloride to yield 121.5 mg (50 % yield) of pure product. M+Li 511

TABLE 8




5

Entry

R


R₁

1


-CH₃

10


2

-CH₃


3

-CH(CH₃)₂

4

-CH(CH₃)₂

5

-C(CH₃)₃

15

TABLE 8 (Cont'd.)












Entry	R	R ₁
5		
6		-CH ₃
7		-CH ₃
8		-CH ₃
9		-CH ₃
10		-CH ₃
11		-CH ₃
12		-CH ₃
13		-CH ₃
14		-CH ₃

TABLE 8 (Cont'd)

15



16




5

Example 18

Following generally the procedure set forth in Example 16, the compounds shown in Table 9 were prepared.

TABLE 9




5

	R ¹	R ^{1'}	R ^{1''}	R
	H	H	H	
	H	H	H	
10	H	CH ₃	H	
	H	CH ₃	CH ₃	
	H	H	CO ₂ CH ₃	
	H	H	H	
	H	H	H	
15	H	H	CONH ₂	Cbz
	H	H	CONH ₂	2-quinolinylcarbonyl
	H	H	CONH ₂	2-benzofurancarbonyl

Example 19

The procedure set forth below in Example 23 was generally utilized to prepare the compounds shown in Table

5 10.

TABLE 10

10

15

20

R	R'	X
R=H	R'=H	X=H
R=Me	R'=Me	X=H
R=H	R'=Me	X=H
R=Me	R'=Me	X=F
R=H	R'=Me	X=F
R=Cbz	R'=Me	X=H
R=H	R'=Bz	X=H
R+R'= pyrrole*		X=H


25

* lle in place of t-butylglycine

Example 20

This example illustrates preparation of compounds wherein R⁴ and R⁵ together with N, forms a heterocycloalkyl radical.

a) Pyrrolidine carbamoyl chloride.




10

A stirring solution of triphosgene (27.78g, 0.103 mol) in 40 mL toluene was cooled to -20 °C in an ice/salt bath under a blanket of nitrogen and treated with a solution of N-methylmorpholine (27.3 g, 0.27 mol) in 20 mL of toluene dropwise over 1h. This solution was then treated with a solution of pyrrolidine (19.8 g, 0.27 mol) in 30 mL of toluene over a period of 30 m. The solution was allowed to warm to room temperature, filtered and the filtrate concentrated in vacuo to give an oil that was purified by vacuum distillation through a 12" Vigeraux column to give 20.7g, 56%, bp 58 °C @ 0.6 mm, of pure product.

20

b) Butanediamide, N¹[3-[[[(4-fluorophenyl)methyl]](1-pyrrolidinylcarbonyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-[1S[1R*(R*),2S*]]-


5



A stirring solution of [1S-[1R*(R*),2S*]]-N¹-[3-
 10 [[(4-fluorophenyl)methyl]amino]-2-hydroxy-1-
 (phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-
 butanediamide (1.08 g, 1.91 mmol) (prepared as in Example
 31) in 7 mL of anhydrous DMF was treated with pyrrolidine
 carbamoyl chloride (260 mg, 1.95 mmol),
 4-dimethylaminopyridine (15 mg), and N-methylmorpholine (380
 15 mg, 3.76 mmol). The solution was stirred at room
 temperature for 3h and then concentrated *in vacuo* to give a
 semi-solid that was dissolved in methanol/water ca. 2:1. A
 solid formed from this solid that was isolated by filtration
 on a Büchner funnel and washed with water, 5% aq. citric
 20 acid and water and air dried to give 130 mg of pure product,
 TLC on SiO₂ eluting with 7% methanol in ethyl acetate showed
 one spot with R_f=0.64, 11%.

c) Butanediamide, N¹[3-[[[(4-fluorophenyl)methyl]](4-morpholinylcarbonyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-[1S[1R*(R*),2S*]]-

5



To a stirring solution of [1S-[1R*(R*),2S*]]-N¹-[3-[[[(4-fluorophenyl)methyl]amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-butanediamide (520 mg, 0.922 mmol), triethylamine (172 mg, 1.70 mmol), 4-dimethylaminopyridine (50 mg), and morpholino carbamoyl chloride (157.3 mg, 1.05 mmol) in 5 mL of chloroform. The initially heterogeneous mixture was heated to reflux for 6 h. The solution was then diluted with additional chloroform, poured into a separatory funnel and washed with 1N KHSO₄, sat. aq. NaHCO₃, dried over anhyd. MgSO₄, filtered, and concentrated in vacuo to give a white solid that was purified by column chromatography on SiO₂ eluting with ethanol/ethyl acetate to give 380 mg, 61%, of pure product.


Example 21

This example illustrates preparation of compounds wherein R⁴ and R⁵ are both other than H.

5

Butanediamide, N¹[3-[[[(diethylamino)carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)-propyl]-2-[(2-quinolinylcarbonyl) amino]-[1S-

10



To a stirring solution of [1S-[1R*(R*),2S*]]-N¹-[3-(methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-
15 [(2-quinolinylcarbonyl)amino-butane diamide] (119 mg, 0.21 mmol) triethylamine (59 mg, 0.58 mmol), 4-dimethylamino-pyridine (9 mg), and diethyl carbamoyl chloride (157.3 mg, 1.05 mmol) in 4 mL of chloroform. The mixture was kept at room temperature for 26 h. The solution was then diluted
20 with additional chloroform, poured into a separatory funnel and washed with 1N KHSO₄, sat. aq. NaHCO₃, dried over anhyd. MgSO₄, filtered, and concentrated in vacuo to give a white solid that was purified by column chromatography on SiO₂ eluting with methanol/CH₂Cl₂ to give 20 mg, 15%, of pure
25 product.

Example 22

Following the procedures set forth in Examples 20 and 21, the compounds listed in Table 11 were prepared.

5

TABLE 11




10	 <chem>Q-ASN-NH-CH(Ph)-CH(OH)-CH2-N(R3)-C(=O)-X(R4)(R5)</chem>	R_3	$\begin{array}{c} X-R_4 \\ \\ R_5 \end{array}$
15	$-\text{CH}_2\text{CH}(\text{CH}_3)_2$ $-\text{CH}_2\text{CH}(\text{CH}_3)_2$ $-\text{CH}_2\text{CH}(\text{CH}_3)_2$	$-\text{N}(\text{CH}_3)_2$ $-\text{N}(\text{CH}_2\text{CH}_3)_2$ $-\text{N}(\text{CH}(\text{CH}_3)_2)_2$	
20	$-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$ $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$ $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$	$-\text{N}(\text{CH}_3)_2$ $-\text{N}(\text{CH}_2\text{CH}_3)_2$	
25	$-\text{CH}_2-$ 	$-\text{N}(\text{CH}_3)_2$ $-\text{N}(\text{CH}_2\text{CH}_3)_2$	

TABLE 11 (Cont'd.)














5	R_3	$\begin{array}{c} X-R_4 \\ \\ R^5 \end{array}$
10	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	
10	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	
15	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	$\text{N}(\text{CH}_3) (\text{t-Bu})$
15	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	
15	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	
20	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	
20	$-\text{CH}_2-\text{C}_6\text{H}_4-\text{F}$	

TABLE 11 (Cont'd.)

5

 R_3
$$\begin{array}{c} X-R_4 \\ | \\ R^5 \end{array}$$

10



15

Example 23A

This example illustrates preparation of compounds of Formula II wherein R¹ is an alkyl group other than an alkyl group of a naturally occurring amino acid side chain. In particular R¹ is a t-butyl group.

5

Part A:

3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino-2(R)-hydroxy-1(S)-[N-(benzyloxycarbonyl)(phenylmethyl)propyl amine] (4.7 gm, 9.7 mmol) was combined with 10% Pd on carbon (200 mg) and conc. HCl (3 mL) in ethanol (35 mL) and hydrogenated at 50 psi of hydrogen for 2.5 h. The reaction mixture was filtered through diatomaceous earth and concentrated on a rotary evaporator to a yellow hygroscopic solid; 3.7 gm, 100%.

15

Part B:

Butaneamide, 2-[(phenylmethyloxycarbonyl)amino]-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl-3,3-dimethyl-1S-[1R*(R*),2S*]]-

20

N-Cbz-L-tert-leucine (172 mg, 0.65 mmol) and N-hydroxybenzotriazole (100 mg, 0.65 mmol) in DMF (3 mL) was cooled to 0°C and EDC (115 mg, 0.60 mmol) added. After 45 min the amine from Part A (193 mg, 0.50 mmol) and N-methylmorpholine (60 uL, 0.55 mmol) were added. The reaction was stirred at ambient temperature for 18 h and poured into a solution of 50% saturated NaHCO₃ (25 mL). The solid was collected by suction filtration, washed with water and dried in-vacuo. The solid was chromatographed on SiO₂ using 2% MeOH in CH₂Cl₂. The appropriate fractions were pooled and concentrated to afford a white solid; 220 mg, MH⁺ 597, TLC (SiO₂ 2%MeOH/CH₂Cl₂) R_f = .2. CHN requires: C, 68.42, H, 8.78, N, 9.39; found: C, 68.03, H, 8.83, N, 9.33.

35

Part C:

Butaneamide, 2-amino-N-[3-[[[(1,1-
dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-2-
hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*),
5 2S*]-

The product from Part B (570 mg, 0.95 mmol) and 4%
Pd on carbon (150 mg) in ethanol (30 mL) was hydrogenated at
5 psi for 2.75 h. The reaction mixture was filtered through
diatomaceous earth and concentrated on a rotary evaporator
10 to an oil; 438 mg, 100%.

Part D:

Butaneamide, 2-(acetyl amino)-N-[3-[[[(1,1-
dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-2-
15 hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*),
2S*]-

The product from Part C (206 mg, 0.41 mmol) and N-
methylmorpholine (45 uL, 0.41 mmol) were dissolved in CH₂Cl₂
(2.5 mL) and cooled to 0 C. Acetic anhydride (39 uL, 0.41
20 mmol) was then added and the reaction stirred 30 min at 0 C,
then allowed to warm to ambient temperature and stir for 30
min. The solvent was removed on a rotary evaporator and the
residue dissolved in ethanol (2 mL). The ethanolic solution
was slowly poured into 50 % saturated NaHCO₃ (20 mL) and
25 stirred vigorously. The solid was collected by suction
filtration and washed with water, 5% citric acid, and again
with water; 157 mg, 75%. CHN / 1.5 H₂O requires: C 63.24, H,
9.67, N, 10.54; found:
C, 63.40, H, 9.41, N, 10.39.

30

Part D2:

Butaneamide, 2-[(2,2-dimethylaminoacetyl)aminol-N-[3-
[[[(1,1-dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-
2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*),
35 2S*]-

To a solution of N,N-dimethylaminoacetic acid (62 mg, 0.6
mmol) and N-hydroxybensotriazole (87 mg, 0.57 mmol) in 2 mL

of DMF at 0° C was added EDC (109 mg, 0.57 mmol). The reaction mixture was stirred for 1 hour and then (2R, 3S)-3(L-tert-butylglyciny) amido-1-isoamyl 1(tert-butylcarboamoyl) amino-4-phenyl-2-butanol (231 mg, 0.5 mmol) was added. The reaction was stirred at room temperature for 16 hours and then concentrated in vacuo. The residue was chromatographed on silica (50 gm) using 5% ethanol in dichloromethane. The butaneamide, 2-[(2,2-dimethylaminoacetyl)amino]-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]- was obtained as a white solid Anal. Calc'd for C₃₀H₅₃N₅O₄.0.3 H₂O: C, 65.13; H, 9.77; N, 12.66. Found: C, 65.10; H, 9.79; N, 12.52.

15

Example 23B

The compound of Example 23A, Part D2, was prepared by an alternate method as described below.

20

To a solution of butaneamide, 2-amino-N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]- (694 mg, 1.5 mmol) and N-methylmorpholine (175 uL, 1.6 mmol) in 5 mL of dichloromethane was added bromoacetyl chloride (132 uL, 1.6 mmol). After 20 minutes at room temperature the solvent was removed on a rotary evaporator and the residue chromatographed on silica (50 gm) using 2% methanol in dichloromethane. The product was obtained as a white solid Anal. Calc'd for C₂₈H₄₇N₄O₄Br.H₂O: C, 55.90; H, 8.21; N, 9.31. Found: C, 55.67; H, 7.90; N, 9.08.

30

To a solution of butaneamide, 2-[(2-bromoacetyl)amino]-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]- (582 mg, 1.0 mmol) and N,N-diisopropylethylamine (260 mL², 1.5 mmol) in 10 mL of dichloromethane was added N,N-dimethylamine hydrochloride (122 mg 1.5 mmol). The

35

reaction was stirred at room temperature for 16 hours, then concentrated on a rotary evaporator. The residue was chromatographed on silica (50 gm) using 2.5% methanol in dichloromethane. The butaneamide, 2-[(2,2-

5 dimethylaminoacetyl)amino]-N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]- was obtained as a white solid Anal Calc'd for $C_{30}H_{53}N_5O_4 \cdot 0.22 CH_2Cl_2$: C, 64.00; H, 9.50; N, 12.35. Found:
10 C, 63.97; H, 9.51; N, 12.24.

Butaneamide, 2-amino-N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]-
15 was also capped, generally utilizing the procedures shown in Examples 23A and 23B, with the acyl groups shown in Table 12.

TABLE 12

5	Acyl Group (R)	
	benzyloxycarbonyl	
	<u>tert</u> -butoxycarbonyl	
	acetyl	
	2-quinoylcarbonyl	
10	phenoxyacetyl	
	benzoyl	
	methyloxaloyl	
	pivaloyl	
	trifluoracetyl	
15	bromoacetyl	
	hydroxyacetyl	
	morpholinylacetyl	
	N-benzylaminoacetyl	
	N-phenylaminoacetyl	
20	N-benzyl-N-methylaminoacetyl	
	N-methyl-N-(2-hydroxyethyl) aminoacetyl	
	N-methylcarbamoyl	
	3-methylbutyryl	
	N-isobutylcarbamoyl	
25	succinoyl (3-carboxypropionyl)	
	carbamoyl	
	N-isobutylamino acetyl	
	N,N-diethylamino acetyl	
	N-(2-methoxyethyl) aminoacetyl	
30	N-(S- α -methylbenzyl) aminoacetyl	
	N-(R- α -methylbenzyl) amino acetyl	
	N-(S-2-tetralin) amino acetyl	
	N-(R-2-teralin) amino acetyl	
	N-pyrrolidinylacetyl	
35	N-methyl-N-(2-pyridylethyl) amino acetyl	
	N-tetrahydroisoquinolylaminoacetyl	
	N-p-methoxybenzylamino acetyl	

Example 24A

The procedure described below illustrates preparation of compounds of Formula III.

- 5 Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(2-phenylethylsulfonyl)-, [1S-[1R*(R*),2S*]] and its diastereomer.

10 Part A:

- A solution of methyl methacrylate (7.25 g, 72.5 mmol) and phenethyl mercaptan (10.0 g, 72.5 mmol) in 100 mL of methanol was cooled in an ice bath and treated with sodium methoxide (100 mg, 1.85 mmol). The solution was stirred under nitrogen for 3 h and then concentrated in vacuo to give an oil that was taken up in ether and washed with 1 N aqueous potassium hydrogen sulfate, saturated aqueous sodium chloride, dried over anhydrous magnesium sulfate, filtered and concentrated to give 16.83 g, 97.5% of methyl 2-(R,S)-methyl-4-thia-6-phenyl hexanoate as an oil. TLC on SiO₂ eluting with 20:1 hexane:ethyl acetate (v:v) R_f=0.41.

Part B:

- 25 A solution of methyl 2-(R,S)-methyl-4-thia-6-phenyl hexanoate (4.00 g, 16.8 mmol) in 100 mL of dichloromethane was stirred at room temperature and treated portion wise with meta-chloroperoxybenzoic acid (7.38 g, 39.2 mmol) over approximately 40 m. The solution was stirred at room temperature for 16 h and then filtered and the filtrate washed with saturated aqueous sodium bicarbonate, 1N sodium hydroxide, saturated aqueous sodium chloride, dried over anhydrous magnesium sulfate, filtered, and concentrated to give 4.50 g, 99% of desired sulfone.
- 35 The unpurified sulfone was dissolved in 100 mL of tetrahydrofuran and treated with a solution of lithium hydroxide (1.04 g, 24.5 mmol) in 40 mL of water. The solution was stirred at room temperature for 2 m and then

concentrated in *vacuo*. The residue was then acidified with 1N aqueous potassium hydrogen sulfate to pH=1 and then extracted three times with ethyl acetate. The combined ethyl acetate solution was washed with saturated aqueous sodium chloride, dried over anhydrous magnesium sulfate, filtered and concentrated to give a white solid. The solid was taken up in boiling ethyl acetate/hexane and allowed to stand undisturbed whereupon white needles formed that were isolated by filtration and air dried to give 3.38 g, 79% of 2-(R,S)-methyl-3(β -phenethylsulfonyl)-propionic acid, mp 91-93°C.

Part C:

A solution of 2-(R,S)-methyl-3(β -phenethylsulfonyl)-propionic acid (166.1 mg, 0.65 mmol), N-hydroxybenzotriazole (HOBT) (146.9 mg, 0.97 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (145.8 mg, 0.75 mmol) in 4 mL of anhydrous dimethylformamide (DMF) cooled to 0°C and stirred under nitrogen for 0.5 h. This solution was then treated with 3-[[[(dimethylethyl)amino]carbonyl](3-methylbutyl)amino-2(R)-hydroxy-1(S)-(phenylmethyl)propyl amine (201.9 mg, 0.59 mmol) and stirred at room temperature for 16 h. The solution was poured into 30 mL of 60% saturated aqueous sodium bicarbonate solution. The aqueous solution was then decanted from the organic residue. The organic residue was taken up in dichloromethane and washed with 10% aqueous citric acid, brine, dried over anhydrous magnesium sulfate, filtered and concentrated to give 110.0 mg, 32% of (2R,3S)-3-[N-2-(R)-methyl-3-(β -phenethylsulfonyl)propionyl]amido-1-isoamyl-1-(tert-butylcarbonyl)amino-4-phenyl-2-butanol and (2R,3S)-3-[N-2-(S)-methyl-3-(β -phenethylsulfonyl)propionyl]amido-1-isoamyl-1-(tert-butylcarbonyl)amino-4-phenyl-2-butanol, FAB mass spectrum (MH⁺) =588. Flash chromatography of the mixture on silica gel eluting with 1:1 hexane:ethyl acetate afforded the separated diastereomers.

Example 24B

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)- [1S-[1R*(R*), 2S*]], and its diastereomer.

Part A:

A solution of methyl 2-(bromomethyl)-acrylate (26.4 g, 0.148 mol) in 100 mL of methanol was treated with sodium methanesulfinat (15.1 g, 0.148 mol) portion wise over 10 m at room temperature. The solution was then stirred at room temperature for a period of 1.25 h and the solution concentrated in vacuo. The residue was then taken up in water and extracted four times with ethyl acetate. The combined ethyl acetate solution was washed with saturated sodium chloride, dried over anhydrous magnesium sulfate, filtered and concentrated to give a white solid, 20.7 g which was taken up in boiling acetone/methyl tert-butyl ether and allowed to stand whereupon crystals of pure methyl 2-(methylsulfonylmethyl) acrylate 18.0 g, 68% formed, mp 65-68 °C.

Part B:

A solution of methyl 2-(methylsulfonylmethyl) acrylate (970 mg, 5.44 mmol) in 15 mL of tetrahydrofuran was treated with a solution of lithium hydroxide (270 mg, 6.4 mmol) in 7 mL of water. The solution was stirred at room temperature for 5 m and then acidified to pH=1 with 1 N aqueous potassium hydrogen sulfate and the solution extracted three times with ethyl acetate. The combined ethyl acetate solution was dried over anhydrous magnesium sulfate, filtered, and concentrated to give 793 mg, 89% of 2-(methylsulfonylmethyl) acrylic acid, mp 147-149 °C.

Part C:

A solution of 2-(methylsulfonylmethyl) acrylic acid (700 mg, 4.26 mmol) in 20 mL of methanol was charged into a Fisher-Porter bottle along with 10% palladium on carbon catalyst under a nitrogen atmosphere. The reaction vessel was sealed and flushed five times with nitrogen and then five times with hydrogen. The pressure was maintained at 50 psig for 16 h and then the hydrogen was replaced with nitrogen and the solution filtered through a pad of celite to remove the catalyst and the filtrate concentrated in *vacuo* to give 682 mg 96% of 2-(R,S)-methyl-3-methylsulfonyl propionic acid.

Part D:

A solution of 2-(R,S)-methyl-3(methylsulfonyl) propionic acid (263.5 mg, 1.585 mmol), N-hydroxybenzotriazole (HOBT) (322.2 mg, 2.13 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (339.1 mg, 1.74 mmol) in 4 mL of anhydrous dimethylformamide (DMF) cooled to 0°C and stirred under nitrogen for 0.5 h. This solution was then treated with 3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2(R)-hydroxy-1(S)-(phenylmethyl)propyl amine (543.5 mg, 1.58 mmol) and stirred at room temperature for 16 h. The solution was poured into 60 mL of 60% saturated aqueous sodium bicarbonate solution. The aqueous solution was then decanted from the organic residue. The organic residue was taken up in dichloromethane and washed with 10% aqueous citric acid, brine, dried over anhydrous magnesium sulfate, filtered and concentrated to give 471.8 mg, 60% of Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]- and its diastereomer.

Example 25Preparation of Sulfone Inhibitors From L-(+)-S-acetyl-β-mercaptoisobutyric Acid5 Part A:

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-S-acetyl)-[1S-[1R*),2S*]]-.

10

A round-bottomed flask was charged with (2R,3R)-3-amino-1-isoamyl-1-(tert-butylcarbonyl)amino-4-phenyl-2-butanol (901.5 mg, 2.575 mmol), L-(+)-S-acetyl-β-mercaptoisobutyric acid (164.5 mg, 2.575 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) (339.1 mg, 1.74 mmol), and 10 mL of CH₂Cl₂ and allowed to stir at room temperature for 16 h. The solution was concentrated in vacuo and the residue taken up in ethyl acetate, washed with 1N KHSO₄ sat. aq. NaHCO₃, brine, dried over anhydrous MgSO₄, filtered and concentrated to give an oil that was purified by radial chromatography on SiO₂ eluting with ethyl acetate to give the pure product, 800 mg, 63%.

25 Part B:

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)-propyl]-2-methyl-3-mercapto)-, [1S-[1R*(R*),2S*]]-.

30

A solution of [1S-[1R*(R*),2S*]]- N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-S-acetyl)-propanamide (420 mg, 0.85 mmol) in 10 mL of methanol was treated with anhydrous ammonia for ca. 1 m at 0°C. The solution was stirred at that temperature for 16 h and then concentrated in vacuo to give 380 mg, 99%, of the desired product that was used directly in the next step without further purification.

Part C:

~~Propanamide~~, N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)-propyl]-2-methyl-3-S-methyl-, [1S-[1R*(R*),2S*]]-.

5

A solution of [1S-[1R*(R*),2S*]]- N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-mercapto)-propanamide (380 mg, 0.841 mmol) in 10 mL of dry toluene under nitrogen was treated in rapid succession with 1,8-diazabicyclo[5.4.0]undec-7-ene, (DBU), (128.1 mg, 0.841 mmol) and iodomethane (119.0 mg, 0.841 mmol). After 0.5 h at room temperature the reaction was found to be complete and the solution was diluted with ethyl acetate washed with 1N KHSO₄, sat. aq. NaHCO₃, brine. After the solution was dried over anhydrous MgSO₄, filtered and concentrated in vacuo the desired product was obtained as white foam was obtained, 370 mg, 94.5%, that was used directed in the next step.

20

Part D:

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)-propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*),2S*]]-.

25


A solution of [1S-[1R*(R*),2S*]]-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-S-methyl)-propanamide (340 mg, 0.73 mmol) and sodium perborate (500 mg, 3.25 mmol) in 30 mL of glacial acetic acid was warmed to 55°C for 16 h. The solution was concentrated in vacuo and then the residue taken up in ethyl acetate, washed with water, sat. aq. NaHCO₃, brine, dried over anhydrous MgSO₄, filtered and concentrated to give the desired product as a white solid, 350 mg, 96%.

35

Example 26

The compounds shown in Tables 13A and 13B were prepared generally according to the procedure set forth in Examples 24 and 25.

TABLE 13A



R'

CH₃-
 CH₃CH₂-
 CH₃CH₂CH₂-
 PhCH₂CH₂-
 PhCH₂-
 Ph-
 (CH₃)₂CH-
 HOCH₂CH₂-







TABLE 13B



5


R'

R₁

10

 $\begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$
 $\begin{matrix} -\text{CH}(\text{CH}_3)_2 \\ -\text{CH}(\text{CH}_3)_3 \end{matrix}$

15

Example 27

20

Preparation of 2(S)-methyl-3-(methylsulfonyl)propionic
Acid.

25

To a solution of 10g of D-(-)-S-benzoyl-b-mercaptioisobutyric acid t-butyl ester in 20 mL of methanol was bubbled in gaseous ammonia at 0°C. The reaction was allowed to then warm to room temperature, stirred overnight and concentrated under reduced pressure. The resulting mixture of a solid (benzamide) and liquid was filtered to provide 5.21g of a pale oil which then solidified. This was identified as 2(S)-methyl-3-mercaptopropionic acid t-butyl ester.

30

To a solution of 5.21g of 2(S)-methyl-3-mercaptopropionic acid t-butyl ester in 75 mL of toluene at 0°C was added 4.50g of 1,8-diazabicyclo[5.4.0]undec-7-ene and 1.94 mL of methyl iodide. After stirring at room temperature for 2.5 hours, the volatiles were removed, ethyl acetate added, washed with dilute hydrochloric acid, water, brine, dried and concentrated to afford 2.82g of a pale oil, identified as 2(S)-methyl-3-(thiomethyl)propionic acid t-butyl ester.

To a solution of 2.82g of 2(S)-methyl-3-(thiomethyl)propionic acid t-butyl ester in 50 mL of acetic acid was added 5.58g of sodium perborate and the mixture heated to 55°C for 17 hours. The reaction was poured into water, extracted with methylene chloride, washed with aqueous sodium bicarbonate, dried and concentrated to afford 2.68g of 2(S)-methyl-3-(methylsulfonyl)propionic acid t-butyl ester as a white solid.


To 2.68g of 2(S)-methyl-3-(methylsulfonyl)-propionic acid t-butyl ester was added 20 mL of 4N hydrochloric acid/dioxane and the mixture stirred at room temperature for 19 hours. The solvent was removed under reduced pressure to afford 2.18g of crude product, which was recrystallized from ethyl acetate/hexane to yield 1.44g of 2(S)-methyl-3-(methylsulfonyl)propionic acid as white crystals.

Example 28

This example illustrates preparation of compounds of Formula IV wherein t is 1.

5

4-N-benzyl itaconamide.




10

A 500 mL three necked round bottomed flask equipped with a dropping funnel, mechanical stirrer, nitrogen inlet and reflux condenser was charged with itaconic anhydride (33.6g, 0.3 mol) and 150 mL of toluene. This solution was added a solution of benzylamine (32.1g, 0.3 mol) in 50 mL of toluene dropwise over 30 m at room temperature. The solution was stirred at this temperature an additional 3h and then the solid product isolated by filtration on a Büchner funnel. The crude product, 64.6g 98%, was recrystallized from 300 mL of isopropyl alcohol to give after two crops 52.1g, 79% of pure product, mp 149-150 °C.

15

20

2(R)-Methyl 4-N-benzyl succinamide.



25


A large Fisher-Porter bottle was charged with the acid from the above reaction (10.95g, 0.05 mol), rhodium (R,R)-DiPAMP (220mg, 0.291 mmol) and 125 mL of degassed methanol. The solution was then hydrogenated at 40 psig for 16h at room temperature. After the hydrogen uptake ceased, the vessel was opened and the solution

30

concentrated in vacuo to give a yellow solid, 11.05g, 100%. The product was then taken up in absolute ethanol and allowed to stand whereupon crystals of the desired product formed, 7.98g, 72%, mp 127-129 °C $[\alpha]_D^{25}$ = +14.9°


- 5 (c=1.332, EtOH), ^1H nmr (CDCl_3) 300MHz 7.30(m, 5H), 6.80(brs, 1H), 4.41(d, $J=5.8\text{Hz}$, 2H), 2.94(m, 1H), 2.62(dd, $J=8.1, 14.9\text{Hz}$, 1H), 2.33(dd, $J=5.5, 14.9\text{Hz}$, 1H), 1.23(d, $J=7.2\text{Hz}$, 3H).

10 4-N(4-methoxybenzyl)itaconamide.



- A 500 mL three necked round bottomed flask
15 equipped with a dropping funnel, mechanical stirrer, nitrogen inlet and reflux condenser was charged with itaconic anhydride (44.8g, 0.4 mol) and 150 mL of toluene. This solution was added a solution of 4-methoxybenzylamine (54.8g, 0.4 mol) in 50 mL of toluene dropwise over 30 m at
20 room temperature. The solution was stirred at this temperature an additional 2h and then the solid product isolated by filtration on a Büchner funnel. The crude product was recrystallized from ethyl acetate/ethanol to
25 give after two crops 64.8g, 65% of pure product, mp 132-134 °C, ^1H nmr (CDCl_3) 300MHz 7.09(d, $J=9.1\text{Hz}$, 2H), 6.90(brt, $J=5.9\text{Hz}$, 1H), 6.74(d, $J=9.1\text{Hz}$, 2H), 6.22(s, 1H), 5.69(s, 1H), 4.24(d, $J=5.9\text{Hz}$, 2H), 3.69(s, 3H), 3.15(s, 2H). ^{13}C nmr (CDCl_3) 170.52, 169.29, 159.24, 135.61, 131.08, 129.37, 128.97, 114.36, 55.72, 43.37, 40.58.


2 (R)-Methyl 4-N(4-methoxybenzyl)succinamide.



5 A large Fisher-Porter bottle was charged with the acid from the above reaction (5.00 g, 0.02 mol), rhodium (R,R)-DiPAMP (110 mg, 0.146 mmol) and 50 mL of degassed methanol. The starting acid was not completely soluble initially, but as the reaction progressed the
10 solution became homogeneous. The solution was then hydrogenated at 40 psig for 16h at room temperature. After the hydrogen uptake ceased, the vessel was opened and the solution concentrated in vacuo to give a yellow solid. The crude product was then taken up in ethyl acetate and washed
15 three times with sat. aq. NaHCO₃ solution. The combined aqueous extracts were acidified to pH=1 with 3 N HCl and then extracted three times with ethyl acetate. The combined ethyl acetate extracts were washed with brine, dried over anhyd. MgSO₄, filtered and concentrated to give
20 the expected product as a white solid, 4.81g, 95%. This material was recrystallized from a mixture of methyl ethyl ketone/hexane to give 3.80g, 75% of pure product, [α]_D @ 25 °C=+11.6° (c=1.572, MeOH). ¹H nmr (CDCl₃) 300MHz 11.9(brs, 1H), 7.18(d, J=9.2Hz, 2H), 6.82(d, J=9.2Hz, 2H), 6.68(brt, J=5.6Hz, 1H), 4.33(d, J=5.6Hz, 2H), 3.77(s, 3H), 2.92(ddq, J=7.9, 5.4, 7.3Hz, 1H), 2.60(dd, J=5.4, 15.0Hz, 1H),
25 2.30(dd, J=7.9, 15.0Hz, 1H), 1.22(d, J=7.3Hz, 3H).

Butanediamide, N'-[3-[[[(1,1-dimethylethyl)amino]-
carbonyl](3-methylbutyl)amino]-2-hydroxy-1-
(phenylmethyl)propyl]-N-4-methoxyphenylmethyl-2-
methyl, [1S-[1R*(2R*),2S*]]-

5




A 50 mL round bottomed flask was charged with
2(R)-methyl 4-N(4-methoxybenzyl)succinamide (588 mg, 2.35
10 mmol), N-hydroxybenzotriazole (511 mg, 3.34 mmol) and 6 mL
of DMF. The solution was cooled to 0° C and treated with
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
(502 mg, 2.62 mmol) for 20 m. A solution of (2R,3S)-3-
amino-1-(3-methylbutyl)-1-[(1,1-
15 dimethylethyl)amino]carbonyl)-4-phenyl-2-butanol (782 mg,
2.24 mmol) in 2 mL of DMF was added and the solution
stirred at room temperature for a period of 24 h. The
solution was concentrated *in vacuo* and poured into 50 mL of
50% sat. aq. NaHCO₃, the aqueous phase was extracted with
20 CH₂Cl₂. The organic phase was washed with 5% citric acid,
NaHCO₃, brine, dried over anhyd. MgSO₄, filtered and
concentrated to give an oil that was purified by radial
chromatography on SiO₂ eluting with hexane/ethyl acetate to
give 790 mg, 59% of pure product as a white foam.

25

Butanediamide, N'-[3-[[[(1,1-dimethylethyl)amino]-
carbonyl](3-methylbutyl)amino]-2-hydroxy-1-
(phenylmethyl)propyl]-N-phenylmethyl-2-methyl, [1S-
[1R*(2R*),2S*]]-

5




A 50 mL round bottomed flask was charged with
2(R)-~~methyl~~ 4-N-(benzyl) succinamide (243 mg, 1.1 mmol), N-
10 hydroxybenzotriazole (213 mg, 1.39 mmol) and 3 mL of DMF.
The solution was cooled to 0° C and treated with 1-(3-
dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (228
mg, 1.17 mmol) for 20 m. A solution of (2R,3S)-3-amino-1-
(3-methylbutyl)-1-[(1,1-dimethylethyl)amino]carbonyl)-4-
15 phenyl-2-butanol (327 mg, 0.95 mmol) in 2 mL of DMF was
added and the solution stirred at room temperature for a
period of 24 h. The solution was concentrated in vacuo and
poured into 50 mL of 50% sat. aq. NaHCO₃, the aqueous phase
was extracted with CH₂Cl₂. The organic phase was washed
20 with 5% citric acid, NaHCO₃, brine, dried over anhyd.
MgSO₄, filtered and concentrated to give an oil that was
purified by flash chromatography on SiO₂ eluting with
hexane/ethyl acetate to give 370 mg, 70% of pure product as
a white foam.

25

Example 29

Following the procedure generally as set forth in Example 28, as well as in Examples 31-45, the compounds shown in Table 14 were prepared.

TABLE 14

R ¹	R ³⁰	R ³¹	R ³²	X'	R ³³	R ³⁴
H	H	H	H	N	H	H
H	H	H	H	O	H	-
H	H	H	H	O	CH ₃	-
CH ₃	H	H	H	N	H	H
CH ₃	H	H	H	O	H	-
H	H	CH ₃	H	N	H	H
H	H	CH ₃	H	O	H	-

TABLE 14 (Cont'd.)

	R1	R30	R31	R32	X'	R33	R34
5	CH ₃	CH ₃	H	H	N	H	H
	CH ₃	CH ₃	H	H	O	H	-
	CH ₃	CH ₃	H	H	O	CH ₂ C ₆ H ₄ OCH ₃	-
10	H	H	CH ₃	CH ₃	N	H	H
	H	H	CH ₃	CH ₃	O	H	-
	H	H	CH ₃	CH ₃	O	CH ₂ C ₆ H ₄ OCH ₃	-
15	CH ₃	H	CH ₃	H	N	H	H
	CH ₃	H	CH ₃	H	N	H	CH ₃
	CH ₃	H	CH ₃	H	N	CH ₃	CH ₃
	CH ₃	H	CH ₃	H	O	H	-
	CH ₃	H	CH ₃	H	N	H	-CH ₂ C ₆ H ₅ OCH ₃
20	OH	H	H	H	N	H	H
	OH	H	H	H	O	H	-
	H	H	OH	H	N	H	H
	H	H	OH	H	O	H	-
25	CH ₂	H	H	H	N	H	H
	CH ₂ C(O)NH ₂	H	H	H	N	H	H
	CH ₂ C(O)NH ₂	H	H	H	O	H	-
30	CH ₂ C(O)NH ₂	H	H	H	O	CH ₃	-
	CH ₂ Ph	H	H	H	N	H	H
	CH ₃	H	CH ₃	CH ₃	N	H	H
35	CH ₃	H	CH ₃	CH ₃	O	H	-
	CH ₃	H	CH ₃	CH ₃	N	H	CH ₃
	CH ₃	H	CH ₃	CH ₃	N	CH ₃	CH ₃

Example 30

Following the procedure generally as set forth
5 in Example 28, as well as Examples 31-45, the compounds
shown in Table 15 were prepared.

TABLE 15







TABLE 15 (Cont'd.)

5

Example 31

10

Preparation of 3(S)-[N-(2-quinolinylcarbonyl)-L-asparaginyl]amino-2(R)-hydroxy-4-phenylbutylamine, N-(3-methylbutyl).

15 Part A:

Preparation of N-3(S)-(Benzyloxycarbonyl)-amino-2(R)-hydroxy-4-phenylbutylamine, N-(3-methylbutyl). A solution of 20g (67 mmol) of N-benzyloxycarbonyl-3(S)-amino-1,2-(S)-epoxy-4-phenylbutane in 140 mL of isopropyl alcohol was treated with 83g (952 mmol) of isoamylamine and refluxed for one hour. The solution was cooled, concentrated, hexane added and the resulting solid filtered to afford 22.4g of the desired product.

25 Part B:

Preparation of N-3(S)-(Benzyloxycarbonyl)-amino-2(R)-hydroxy-4-phenylbutylamine, N-(3-methylbutyl)-N-(t-butyloxycarbonyl). To a solution of 22.4g (58.3 mmol) of product from Part A above, 6.48g (64.1 mmol) of triethylamine and 150 mg of N,N-dimethyl-4-aminopyridine in 200 mL of tetrahydrofuran at 0°C was added 12.7g (58.3 mmol) of di-t-butylpyrocarbonate in 10 mL of THF. After

30

3.5 hours at room temperature, the volatiles were removed, ethyl acetate added and washed with 5% citric acid, sat'd ~~NaHCO~~₃, dried and concentrated to afford 30g of crude product. Chromatography on silica gel using 20% ethyl acetate/hexane afforded 22.5g (79%) of the desired product.

Part C:

Preparation of N-3(S)-[N-benzyloxycarbonyl-L-asparaginy]amino-2(R)-hydroxy-4-phenylbutylamine, N-(3-methylbutyl)-N-(t-butyloxycarbonyl). A solution of 22.5g of product from Part B above in 200 mL of ethanol was hydrogenated over 5.9g of 10% palladium-on-carbon under 50 psig hydrogen for one hour. The catalyst was filtered and the solvent removed under reduced pressure to afford 15.7g of free amine. This was dissolved in 130 mL of DMF and 4.54g (44.9 mmol) of N-methylmorpholine was added to a mixture of 13.3g (49.9 mmol) N-benzyloxy-carbonyl-L-asparagine, 11.5g (74.9 mmol) of N-hydroxybenzotriazole and 10.5g (54.9 mmol) of EDCI in 120 mL of DMF at 0°C, which had been preactivated for one hour prior to the addition. The mixture was stirred for 2 hours at 0°C and then for 12 hours at room temperature. The reaction was poured into 1L of sat'd aqueous sodium bicarbonate, the solid collected, dissolved in ethyl acetate, washed with water, sat'd sodium bicarbonate, 5% citric acid and brine, dried and concentrated to afford 16.7g of the desired product.

Part D:


Preparation of N-3(S)-[N-(2-quinolinylcarbonyl)-L-asparaginy]amino-2(R)-hydroxy-4-phenylbutylamine, N-(3-methylbutyl)-N-(t-butyloxycarbonyl). A solution of 16.7g (28.0 mmol) of product from Part C in 250 mL of methanol was hydrogenated over 6.0g of 10% palladium-on-carbon and under 50 psig hydrogen for one hour. The catalyst was filtered and the solution concentrated to afford 10.0g of free amine. This was dissolved in 100 mL of methylene chloride, 4.35g (43 mmol) of N-methylmorpholine was added followed by 5.53g (20.5 mmol) of quinoline-2-carboxylic

acid, N-hydroxysuccinimide ester. This was stirred at room temperature overnight, the solvent removed, ethyl acetate added and washed with 5% citric acid, sat'd sodium bicarbonate, brine, dried and concentrated to afford 14g of crude product. Recrystallization from ethyl acetate and hexane afforded 10.5g (83%) of desired product.

Part E:

Preparation of N-3(S)-[N-(2-quinolinyl-carbonyl)-L-asparaginy]amino-2(R)-hydroxy-4-phenylbutylamine, N-(3-methylbutyl). To 80 mL of 4N hydrochloric acid in dioxane was added 9.17g (14.8 mmol) of product from Part D above. After one hour, the product becomes gummy. The solvents were removed, diethyl ether added and removed and the residue dissolved in 20 mL of methanol. This solution was added to 400 mL of sat'd aqueous sodium bicarbonate, the solids collected, washed with acetone and hexane and dried in vacuo over P₂O₅ to afford 4.75g of the desired product.

Example 32A



Preparation of Benzyl 2,2,3(R)-trimethylsuccinate

Part A:

Preparation of Methyl (S)-lactate, 2-methoxy-2-propyl ether. To a mixture of methyl(s)-(-)-lactate (13.2g, 100 mmol) and, 2-methoxypropene (21.6g, 300 mmol) in CH₂Cl₂ (150 ml) was added POCl₃ (7 drops) at r.t. and the resulting mixture was stirred at this temperature for 16 hours. After the addition of Et₃N (10 drops), the solvents were removed in vacuo to give 20.0g of (98%) desired product.

Part B:

Preparation of 2(S)-hydroxypropanal, 2-methoxy-2-propyl ether. To a solution of compound from Part A (20.0g) in CH₂Cl₂ (100 ml) was added DIBAL (65 ml of 1.5M solution in toluene, 97.5 mmol) dropwise at -78°C for 45 min., then stirring was continued at the temperature for another 45 min. To this cold solution was added MeOH (20 ml), saturated NaCl solution (10 ml) and allowed the reaction mixture to warm up to r.t. and diluted with ether (200 ml), MgSO₄ (150g) was added and stirred for another 2 h. The mixture was filtered and the solid was washed twice with ether. The combined filtrates were rotavaped to afford 11.2g (78%) of the desired aldehyde.

15 Part C:

Preparation of 2(S)-hydroxy-cis-3-butene, 2-methoxy-2-propyl ether. To a suspension of ethyltriphenylphosphonium bromide (28g, 75.5 mmol) in THF (125 ml) was added KN (TMS)₂ (15.7g, 95%, 75 mmol) in portions at 0°C and stirred for 1 h at the temperature. This red reaction mixture was cooled to -78°C and to this was added a solution of aldehyde from Part B (11g, 75 mmol) in THF (25 ml). After the addition was completed, the resulting reaction mixture was allowed to warm up to r.t. and stirred for 16 h. To this mixture was added saturated NH₄Cl (7.5 ml) and filtered through a pad of celite with a thin layer of silica gel on the top. The solid was washed twice with ether. The combined filtrates were concentrated in vacuo to afford 11.5g of crude product. The purification of crude product by flash chromatography (silica gel, 10:1 Hexanes/EtoAc) affording 8.2g (69%) pure alkene.

30 Part D:

35 Preparation of 2(S)-hydroxy-cis-3-butene. A mixture of alkene from Part C (8.2g) and 30% aqueous acetic acid (25 ml) was stirred at r.t. for 1 hour. To this mixture was added NaHCO₃ slowly to the pH ~ 7, then

extracted with ether (10 ml x 5). The combined ether solutions were dried (Na_2SO_4) and filtered. The filtrate was distilled to remove the ether to give 2.85g (64%) pure alcohol, $m/e=87(\text{M}+\text{H})$.

5

Part E:

Preparation of 2,2,3-trimethyl-hex-(trans)-4-enoic acid. To a mixture of alcohol from Part D (2.5g, 29 mmol) and pyridine (2.5 ml) in CH_2Cl_2 (60 ml) was added isobutyryl chloride (3.1g, 29 mmol) slowly at 0°C . The resulting mixture was stirred at r.t. for 2 hours then washed with H_2O (30 ml x 2) and sat. NaCl (25 ml). The combined organic phases were dried (Na_2SO_4), concentrated to afford 4.2g (93%) ester 2(S)-hydroxy-cis-3-butenyl isobutyrate. This ester was dissolved in THF (10 ml) and was added to a 1.0M LDA soln. (13.5 ml of 2.0M LDA solution in THF and 13.5 ml of THF) slowly at -78°C . The resulting mixture was allowed to warm up to r.t. and stirred for 2 h and diluted with 5% NaOH (40 ml). The organic phase was separated, the aqueous phase was washed with Et_2O (10 ml). The aqueous solution was collected and acidified with 6N HCl to pH ~ 3. The mixture was extracted with ether (30 ml x 3). The combined ether layers were washed with sat. NaCl (25 ml), dried (Na_2SO_4) and concentrated to afford 2.5g (60%) of desired acid, $m/e=157(\text{M}+\text{H})$.

25

Part F:

Preparation of benzyl 2,2,3(S)-trimethyl-trans-4-hexenoate. A mixture of acid from Part E (2.5g, 16 mmol), BnBr (2.7g, 15.8 mmol), K_2CO_3 (2.2g, 16 mmol), NaI (2.4g) in acetone (20 ml) was heated at 75°C (oil bath) for 16 h. The acetone was stripped off and the residue was dissolved in H_2O (25 ml) and ether (35 ml). The ether layer was separated, dried (Na_2SO_4) and concentrated to afford 3.7g (95%) of benzyl ester, $m/e=247(\text{M}+\text{H})$.

35

Part G:

Preparation of benzyl 2,2,3(R)-trimethylsuccinate. To a well-stirred mixture of KMnO_4 (5.4g, 34, 2 mmol), H_2O (34 ml), CH_2Cl_2 (6 ml) and

5 benzyltriethylammonium chloride (200 mg) was added a solution of ester from Part F (2.1g, 8.54 mmol) and acetic acid (6 ml) in CH_2Cl_2 (28 ml) slowly at 0°C . The resulting mixture was stirred at the temperature for 2 h then r.t. for 16 h. The mixture was cooled in an ice-water bath, to

10 this was added 6N HCl (3 ml) and solid NaHSO_3 in portions until the red color disappeared. The clear solution was extracted with CH_2Cl_2 (30 ml x 3). The combined extracts were washed with sat. NaCl solution, dried (Na_2SO_4) and concentrated to give an oil. This oil was dissolved in

15 Et_2O (50 ml) and to this was added sat. NaHCO_3 (50 ml). The aqueous layer was separated and acidified with 6N HCl to pH ~ 3 then extracted with Et_2O (30 ml x 3). The combined extracts were washed with sat. NaCl solution (15 ml), dried (Na_2SO_4) and concentrated to afford 725 mg (34%)

20 of desired acid, benzyl 2,2,3(R)-trimethylsuccinate, $m/e=251(\text{M}+\text{H})$.

Example 32BPart A:

25 Preparation of Butanediamide, N¹[3-[[[(1,1-dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-2-hydroxy-1-(phenylmethyl)propyl]-2,3,3-trimethyl-1S-, [1R*(2S*),2S*]]-

30 To a well-stirred solution of benzyl 2,2,3(R)-trimethylsuccinate (225 mg, 0.9 mmol) in DMF (1.0 ml) was added HOBt (230 mg, 1.5 mmol). The clear reaction mixture was then cooled to 0°C , to this was added EDC (210 mg, 1.1 mmol) and stirred for 1 h at the temperature. To this cold

35 mixture was added a powder of (350 mg, 1.0 mmol) and DMF (0.5 ml). The resulting reaction mixture was stirred for 2 h at 0°C and 16 h at r.t. After the removal of DMF ($\leq 40^\circ\text{C}$), a solution of 60% sat. NaHCO_3 (10 ml) was added.

This mixture was extracted with EtOAc (10 ml x 2). The extracts were combined and washed with sat. NaHCO₃ (10 ml x 2), 5% citric acid (10 ml x 2), H₂O (10 ml), sat. NaCl (10 ml) and dried (Na₂SO₄) then concentrated to afford 512 mg (98%) of desired product Butanoic Acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-2,2,3-trimethyl-4-oxo-, [1S-[1R*(3S*),2S*]]-benzyl ester as a white solid, m/e=582 (M+H).

Part B:

A mixture of benzyl ester from Part A (480 mg, 0.825 mmol), 10% Pd/C (450 mg) in MeOH (25 ml) was hydrogenated (H₂, 50 psi) for 1/2 h at r.t. The mixture was filtered and the solid was washed with MeOH (10 ml). The collected filtrates were concentrated to afford a crude acid as a white solid. The crude acid was dissolved in Et₂O-EtOAc (10:1, 25 ml) and the solution was washed with sat. NaHCO₃ (25 ml) then 5% NaOH (10 ml). The combined aqueous layers were cooled to 0°C and acidified with concentrated HCl (Co₂) to pH ~ 1 then extracted with Et₂O-EtOAc (10:1, 25 ml x 3). The combined extracts were washed with sat. NaCl (15 ml), dried (Na₂SO₄) and concentrated to afford 307 mg (75.7%) of pure acid Butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-2,2,3-trimethyl-4-oxo-, [1S-[1R*(3S*),2S*]]-, as a white solid, m/e=491 (M+H).


Part C:

Butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-2,2,3-trimethyl-4-oxo-, [1S-[1R*(3S*),2S*]]-, as a white solid, m/e=491 (M+H).

To a well-stirred solution of the acid from Part B (245 mg, 0.5 mmol) in DMF (0.5 ml) was added HOBt (153 mg, 1.0 mmol) and EDC (143 mg, 0.75 mmol) at 0°C. After stirring at 0°C for 2 h, NH₄OH (0.63 ml of 28% NH₄OH, 5

mmol) was added and stirred at 0°C for 2 h, r.t. for 16 h. The removal of DMF ($\leq 40^\circ\text{C}$) gave a white solid. The purification of the crude product by flash chromatography (silica gel, 5% MeOH/CH₂Cl₂) gave 172 mg (70%) of pure
5 amide as a white solid, m/e=491(M+H).

Example 33



Preparation of methyl 2,2-dimethyl-3-methyl succinate, (R) and (S) isomers.

15 Part A:

Preparation of methyl 2,2-dimethyl-3-oxo-butanoate. A 250 ml RB flask equipped with magnetic stir bar and N₂ inlet was charged with 100 ml dry THF and 4.57g (180 mmol) of 95% NaH. The slurry was cooled to -20°C and
20 10g (87 mmol) methyl acetoacetate was added dropwise followed by 11.3 ml (181 mmol) CH₃I. The reaction was stirred at 0°C for 2 hours and let cool to room temperature overnight. The reaction was filtered to remove NaI and diluted with 125 ml Et₂O. The organic phase was washed
25 with 1x100 l 5% brine, dried and concentrated in vacuo to a dark golden oil that was filtered through a 30g plug of silica gel with hexane. Concentration in vacuo yielded 10.05g of desired methyl ester, as a pale yellow oil, suitable for use without further purification.

30

Part B:

Preparation of methyl 2,2-dimethyl-3-oxo-(trifluoromethanesulfonate)-but-3-enoate. A 250 ml RB flask equipped with magnetic stir bar and N₂ inlet was
35 charged with 80 mL by THF and 5.25 ml (37.5 mmol)

diisopropylamine was added. The solution was cooled to -25°C (dry ice/ethylene glycol) and 15 ml (37.5 mmol) of 2.5 M n-BuLi in hexanes was added. After 10 minutes a solution of 5g (35 mmol) **1** in 8 ml dry THF was added. The deep yellow solution was stirred at -20°C for 10 min. then 12.4g N-phenyl bis(trifluoromethanesulfonimide) (35 mmol) was added. The reaction was stirred @ -10°C for 2 hours, concentrated in vacuo and partitioned between ethyl acetate and sat. NaHCO₃. The combined organic phase was washed with NaHCO₃, brine and conc. to an amber oil that was filtered through 60g silica gel plug with 300 mL 5% ethyl acetate/hexane Conc. in vacuo yielded 9.0g light yellow oil that was diluted with 65 ml ethyl acetate and washed with 2x50 ml 5% aq K₂CO₃, 1x10 mL brine, dried over Na₂SO₄ and conc. in vacuo to yield 7.5g (87%) vinyl triflate, (m/e=277 (M+H)) suitable for use without further purification.

Part C:

Preparation of methyl 2,2-dimethyl-3-carboxyl-but-3-enoate. A 250 ml Fisher Porter bottle was charged with 7.5g (27 mmol) of compound prepared in B, 50 ml dry DMF, 360 mg (1.37 mmol) triphenyl phosphine and 155 mg (.69 mmol) Pd(II) (OAc)₂. The reaction mixture was purged twice with N₂ then charged with 30 psi CO. Meanwhile a solution of 20 ml dry DMF and 7.56 ml (54 mmol) NEt₃ was cooled to 0°C to this was added 2.0g (43 mmol) of 99% formic acid. The mixture was swirled and added to the vented Fisher Porter tube. The reaction vessel was recharged to 40 psi of CO and stirred 6 hours @ room temperature. The reaction mixture was concentrated in vacuo and partitioned between 100 mL of ethyl acetate and 75 mL 5% aq K₂CO₃. The aqueous phase was washed with 1x40 mL additional ethyl acetate and then acidified with conc. HCl/ice. The aqueous phase was extracted with 2x70 mL of ethyl acetate and the organics were dried and conc. to yield 3.5g (75%) white crystals, mp 72-75°C, identified as the desired product (m/e=173 (M+H)).

Part D:

Preparation of methyl 2,2-dimethyl-3-methylsuccinate, isomer #1. A steel hydrogenation vessel was charged with 510 mg (3.0 mmol) acrylic acid, from Part C, and 6 mg Ru (acac)₂ (R-BINAP) in 10 ml degassed MeOH. The reaction was hydrogenated at 50 psi/room temperature for 12 hours. The reaction was then filtered through celite and conc. to 500 mg clear oil which was shown to be a 93:7 mixture of isomer #1 and #2, respectively as determined by GC analysis using a 50 M β -cyclodextrin column: 150°C - 15 min. then ramp 2°C/min.; isomer #1, 17.85 min., isomer #2, 18-20 min.

Part E:

Preparation of methyl 2,2-dimethyl-3-methylsuccinate, Isomer #2. A steel hydrogenation vessel was charged with 500 mg (2.9 mmol) acrylic acid, Part C, and 6 mg Ru(OAc) (acac) (S-BINAP) in 10 ml degassed MeOH. The reaction was hydrogenated at 50 psi/room temperature for 10 hours. The reaction was filtered through celite and concentrated in vacuo to yield 490 mg of product as a 1:99 mixture of isomers #1 and #2, respectively, as determined by chiral GC as above.

In a similiar manner, one can use benzyl 2,2-dimethyl-3-oxo-butanoate to prepare benzyl 2,2,3-trimethylsuccinate, R and S isomers.

Example 34

Preparation of 3-[[[(1,1-dimethylethyl)aminolcarbonyl]- (3-methylbutyl)aminol-2(R)-hydroxy-1(S)- (phenylmethyl)propylamine, 1.

Part A:

To a solution of 75.0g (0.226 mol) of N-benzyloxycarbonyl-L-phenylalanine chloromethyl ketone in a mixture of 807 mL of methanol and 807 mL of tetrahydrofuran

at -2°C, was added 13.17g (0.348 mol, 1.54 equiv.) of solid sodium borohydride over one hundred minutes. The solvents were removed under reduced pressure at 40°C and the residue dissolved in ethyl acetate (approx. 1L). The solution was
5 washed sequentially with 1M potassium hydrogen sulfate, saturated sodium bicarbonate and then saturated sodium chloride solution. After drying over anhydrous magnesium sulfate and filtering, the solution was removed under reduced pressure. To the resulting oil was added hexane
10 (approx. 1L) and the mixture warmed to 60°C with swirling. After cooling to room temperature, the solids were collected and washed with 2L of hexane. The resulting solid was recrystallized from hot ethyl acetate and hexane to afford 32.3g (43% yield) of N-benzyloxycarbonyl-3(S)-
15 amino-1-chloro-4-phenyl-2(S)-butanol, mp 150-151°C and $M+Li^+ = 340$.

Part B:

To a solution of 6.52g (0.116 mol, 1.2 equiv.) of
20 potassium hydroxide in 968 mL of absolute ethanol at room temperature, was added 32.3g (0.097 mol) of N-CBZ-3(S)-amino-1-chloro-4-phenyl-2(S)-butanol. After stirring for fifteen minutes the solvent was removed under reduced pressure and the solids dissolved in methylene chloride. After washing with
25 water, drying over magnesium sulfate, filtering and stripping, one obtains 27.9g of a white solid. Recrystallization from hot ethyl acetate and hexane afforded 22.3g (77% yield) of N-benzyloxycarbonyl-3(S)-amino-1,2(S)-epoxy-4-phenylbutane, mp 102-103°C and $MH^+ 298$.

30

Part C:

A solution of N-benzyloxycarbonyl 3(S)-amino-1,2-(S)-epoxy-4-phenylbutane (30.1g, 0.10 mol) and 165mL of isoamylamine in 150 mL of isopropyl alcohol was heated to
35 reflux for 2.5 hours. The solution was cooled to room temperature, concentrated in vacuo and then recrystallized. The product was isolated by filtration and from ethyl

acetate/hexane to afford 31.7g (81%) of N[3(S)-benzyloxycarbonylamino-2(R)-hydroxy-4-phenylbutyl]N-isoamylamine.

5 Part D:

A solution of N[3(S)-benzyloxycarbonylamino-2(R)-hydroxy-4-phenyl butyl], N-isoamylamine in 10 ml of tetrahydrofuran was treated with tert-butylisocyanate (267 mg, 2.70 mmol) at room temperature for 5 minutes. The solvent was removed in vacuo and replaced with ethyl acetate. The ethyl acetate solution was washed with 5% citric acid, water, and brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo to give 1.19g, 97% of N-benzyloxycarbonyl-3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2(R)-hydroxy-1(S)-(phenylmethyl)propylamine, MH⁺ m/z = 470.

Part E:

A solution of (37.3g, 77 mmol) of product from Part D in 100 mL of methanol was hydrogenated over 10% palladium-on-carbon for 4 hours to afford 26.1g of the desired product.

Example 35

Preparation of Butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-, [1S-[1R*, 2S*]]-

30 Part A:

To a solution of 102mg (0.29 mmol) of Example 34 Part E and 70 mg (0.89 mmol) of pyridine in 2 mL of methylene chloride was added 29 mg (0.29 mmol) of succinic anhydride. After 2 hours, ethyl acetate was added and then extracted with saturated NaHCO₃. The aqueous layer was acidified, reextracted with ethyl acetate, washed with saturated brine, dried over magnesium sulfate, filtered and concentrated in vacuo to afford 78 mg (60%) of butanoic

acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino-4-oxo-, [1S-[1R*, 2S*]-.

5 Part B:

his was activated with EDC and N-hydroxybenzotriazole in N,N-dimethylformamide and then reacted with ammonia to generate the desired final compound.

10

Example 36

Part A:

To a solution of 4.60g (24.7 mmol) of trans-diethyl
15 1,2-cyclopropanedicarboxylate in 100 mL of 50:50 v:v tetrahydrofuran/water was added 1.24g (29.6 mmol) of lithium hydroxide. After 17 hours, the tetrahydrofuran was removed in vacuo, the water layer washed with ethyl acetate, acidified with 1N hydrochloric acid and reextracted with ethyl acetate.
20 The organic layer was dried and stripped to afford 2.1g of crude product. After recrystallization from diethyl ether/hexane and then methylene chloride/hexane one obtains 1.1g (28%) of trans-monoethyl 1,2-cyclopropanedicarboxylate, m/e = 159 (M + H).

25

Part B:

To a solution of 297 mg (1.87 mmol) of trans-monoethyl 1,2-cyclopropanedicarboxylate and 429 mg (2.8 mmol) N-hydroxybenzotriazole (HOBT) in 3 mL of anhydrous
30 N,N-dimethylformamide (DMF) at 0°C was added 394 mg (2.0 mmol) of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC). After 30 min. a solution of 591 mg (1.7 mmol) of product from Part A in 2 mL DMF and 171 mg (1.69 mmol) of N-methylmorpholine (NMM) was added. After 2
35 hours at 0°C, the reaction was stirred at RT overnight, poured into water, extracted with ethyl acetate, washed with water, 5% aq. citric acid, sat'd NaHCO₃, brine, dried over anhydride MgSO₄ and stripped to afford 771 mg of crude

product. This was chromatographed on silica gel using 5-20% methanol/methylene chloride to afford 670 mg (80%) of cyclopropane carboxylic acid, 2-[[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]carbonyl]-, ethyl ester; $m/e = 490$ ($M + H$).

Part C:

To a solution of 658 mg (1.32 mmol) of product from Part B in 5 mL of 50:50 THF/water was added 66 mg (1.58 mmol) of lithium hydroxide. After 19 hours, the THF was removed in vacuo, the water washed with ethyl acetate, acidified and reextracted with ethyl acetate. The organic layer was dried and stripped to afford 328 mg (54%) of the corresponding acid, $m/e = 462$ ($M + H$).

Part D:

To a solution of 304 mg (0.66 mmol) of product from Part C, 151 mg (0.99 mmol) HOBT in 2.2 mL DMF at 0°C was added 139 mg (0.73 mmol) EDC. After 30 min. at 0°C, 1.1 mL of conc. aqueous ammonia was added. After stirring at 0°C for 2 hours and RT for 20 hours, the reaction was poured into brine and extracted with ethyl acetate. After washing with sat'd NaHCO_3 , brine, drying and stripping, one obtains 141 mg of crude product. This was chromatographed on silica gel with 1-5% methanol/methylene chloride to afford 40 mg (13%) of the desired final product, $m/e = 561$ ($M + H$).

Example 37

Preparation of trans-but-2-enediamide, N-[3-[[[(1,1-dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-2-hydroxy-1-(phenylmethyl)propyl]-, [1S-[1R*, 2S*].

Part A:

To a solution of 137 mg (0.95 mmol) fumaric acid monoethyl ester in 1 mL of DMF at 0°C was added 183 mg (0.95 mmol) EDCI. After 15 minutes, a solution of 333 mg (0.95 mmol) of the compound of Example 34 Part E in 1 mL DMF was added and the reaction stirred for 14 hours at RT. Ethyl acetate was added and extracted with sat'd brine, 0.2 N HCl, sat'd NaHCO₃, dried and stripped to afford 0.32g of crude product. Chromatography on silica gel using 0-50% ethyl acetate/hexane afforded 0.26g (58%) of but-2-enoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-4-oxo-, [1S-[1R*, 2S*]]-, ethyl ester, m/e = 476 (M + H).

15 Part B:

To a solution of 26.6 mg (0.56 mmol) of product from Part A in 3 mL of 50:50 THF/water was added 34 mg (0.82 mmol) of lithium hydroxide and the reaction stirred at RT for 1 hour. The THF was removed in vacuo, the aqueous layer acidified with 1N HCl and extracted with ethyl acetate. The organic layer was washed with brine, dried and stripped to afford 233 mg (93%) of trans-but-2-enoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-4-oxo-, [1S-[1R*, 2S*]]-, m/e = 448 (M + H).

25 Part C:

To a solution of 225 mg (0.50 mmol) of the product from Part B in 1 mL of DMF was added 95 mg (0.50 mmol) EDC. After 15 minutes at RT, 0.50 mL of conc. aqueous ammonia was added and the reaction stirred for 15 hours. Ethyl acetate was added and washed with 0.2N HCl, brine, dried and stripped to afford 170 mg of crude product. After chromatography on silica gel using 0-40%

methanol/methylene chloride, one obtains 50 mg (22%) of
trans-but-3-enediamide, N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl]-, [1S-[1R*, 2S*]-, m/e =
5 447 (M + H).

Example 38

Preparation of butanediamide, N-[3-[[[(1,1-
10 dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl-2-methyl-, [1S-[1R*(2S*),
2S*1-.

Part A:

15 To a suspension of 24.7g (0.22 mol) of itaconic
anhydride in 100 mL of anhydrous toluene at reflux under a
nitrogen atmosphere was added dropwise over 30 minutes
23.9g (0.22 mol) of benzyl alcohol. The insoluble material
dissolved to provide a homogeneous solution which was
20 refluxed for 1.5 hours. The solution was cooled to RT,
then in an ice bath and the resulting white precipitate
collected by filtration to afford 24.8g (51%) of 4-benzyl
itaconate.

25 Part B:

To a solution of 2.13g (9.5 mmol) of the product
from Part A in 12 mL of methylene chloride at 0°C was added
4.02g (29.1 mmol) of para-methoxybenzyl alcohol, 605 mg
(4.95 mmol) of N,N-dimethyl 4-aminopyridine, 128 mg of N,N-
30 dimethyl 4-aminopyridine hydrochloride salt and then 2.02g
(4.7 mmol) dicyclohexylcarbodiimide (DCC). After stirring
at 0°C for 1 hour and then RT for 2 hours, the precipitate
was collected and discarded. The filtrate was washed with
0.5 N HCl, sat'd NaHCO₃, dried and stripped to afford 4.76g
35 of crude product. This was chromatographed on silica gel
using 0-50% ethyl acetate/hexane to afford 1.24g of pure
4'-methoxybenzyl-4-benzylitaconate.

Part C:

A solution of 1.24g (3.65 mmol) of product from Part B and 20 mg of [(R,R)-Dipamp)cyclooctadienylrhodium] tetrafluoroborate in 30 mL of methanol was thoroughly
5 degassed, flushed with nitrogen and then hydrogen and then stirred under 50 psig of hydrogen for 15 hours. The solution was filtered and stripped, dissolved in methylene chloride and washed with sat'd NaHCO₃, dried and stripped to afford 0.99g of a brown oil. This was then dissolved in
10 40 mL of methylene chloride, 3 mL of trifluoroacetic acid added and the solution stirred at RT for 3.5 hours. Water was added and separated and the organic layer extracted with sat'd NaHCO₃. The aqueous layer was acidified and reextracted with ethyl acetate, separated and the organic
15 layer washed with brine, dried and stripped to afford 320 mg (50%) of 2(R)-methyl-4-benzylsuccinic acid.

Part D:

To a solution of 320 mg (1.44 mmol) of product
20 from Part C and 314 mg (2.05 mmol) HOBt in DMF at 0°C was added 303 mg (1.58 mmol) of EDC. After stirring for 30 minutes, a solution of 467 mg (1.34 mmol) of the product from Example 34, Part E in 4 mL of DMF was added. After
stirring for 1 hour at 0°C and 14 hours at RT, ethyl
25 acetate was added and washed with sat'd NaHCO₃, 5% aqueous citric acid, dried and stripped to afford 0.97g of crude product. This was chromatographed on silica gel using 0-10% ethyl acetate/hexane to afford 420 mg of pure
butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]-
30 carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-3-methyl-4-oxo-, [1S-[1R*(3S*), 2S*]-, benzyl ester.

Part E:

35 A solution of 150 mg (0.27 mmol) of product from Part D in 15 mL of methanol was hydrogenated over 10% palladium on carbon under 50 psig hydrogen for 17 hours. The reaction was filtered and stripped to afford 125 mg

(100%) of butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-3-methyl-4-oxo-, [1S-[1R*(3S*), 2S*]-.

5

Part F:

To a solution of 125 mg (0.27 mmol) of product from Part E and 65 mg (0.42 mmol) of HOBt in 5 mL of DMF at 0°C was added 59 mg (0.31 mmol) of EDC. After 30 min. at 0°C, 1 mL of conc. aqueous ammonia was added. After stirring at 0°C for 2 hours and RT for 15 hours, ethyl acetate was added and washed with sat'd NaHCO₃, 5% aqueous citric acid, dried and stripped to afford 90 mg of crude product. This was recrystallized from ethyl acetate/hexane to afford 40 mg (32%) of pure butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl, [1S-[1R*(2S*), 2S*]-.

20

Example 39

Preparation of butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl, [1S-[1R*(2R*), 2S*]-.

25

Part A:

A solution of 1.41g (4.1 mmol) of 4'-methoxybenzyl 4-benzylitaconate and 25 mg of [(S,S-Dipamp)cyclooctadienylrhodium]tetrafluoroborate in 20 mL of methanol was thoroughly degassed, flushed with nitrogen and then hydrogen and then stirred under 40 psig hydrogen for 72 hours. The solution was filtered and concentrated to provide 1.34g of a brown oil. This was dissolved in 40 mL of methylene chloride and 3 mL of trifluoroacetic acid was added. After stirring for 4 hours, water was added, separated and the organic layer extracted with sat'd NaHCO₃. The aqueous layer was separated, reacidified,

35

extracted with ethyl acetate which was separated, washed with brine, dried and stripped to afford 440 mg of 2(S)-methyl-4-benzylsuccinic acid.

5 Part B:

To a solution of 440 mg (1.98 mmol) of the product from Part A and 437 mg (2.86 mmol) of HoBT in 9 mL of DMF at 0°C was added 427 mg (2.23 mmol) of EDCI. After 30 minutes at 0°C, a solution of 653 mg (1.87 mmol) of the product from Example 34, Part E in 3 mL DMF was added. After 1 hour at 0°C and 15 hours at RT, ethyl acetate was added, extracted with sat'd NaHCO₃, 5% aqueous citric acid, dried and concentrated to afford 0.98g of crude product. Chromatography on silica gel using 0-10% ethyl acetate afforded 610 mg (59%) of pure butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)-amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-3-methyl-4-oxo-, [1S-[1R*(3R*), 2S*], benzyl ester.

20 Part C:

A solution of 310 mg (0.56 mmol) of the product from Part B in 20 mL of methanol was hydrogenated over 20 mg of 10% palladium on carbon under 50 psig hydrogen for 19 hours. The solution was filtered and concentrated to afford 220 mg (85%) of butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)-amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-3-methyl-4-oxo-, [1S-[1R*(3R*), 2S*].

30 Part D:

To a solution of 190 mg (0.41 mmol) of the product from Part C and 90 mg (0.58 mmol) HOBT in 5 mL of DMF at 0°C, was added 88 mg (0.46 mmol) of EDC. After 30 minutes at 0°C, 2 mL of conc. aqueous ammonia was added. After 1 hour at 0°C and 15 hours at RT, ethyl acetate was added, washed with sat'd NaHCO₃, 5% aqueous citric acid, dried and concentrated to afford crude product. Recrystallization from ethyl acetate/hexane afforded 20 mg

(11%) of butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl, [1S-[1R*(2R*), 2S*]].

5

Example 40

Preparation of butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3-methyl-, [1S-[1R*(3S*), 2S*]]-.

10

Part A:

In a similar manner to the procedure used above, p-methoxybenzyl alcohol was reacted with itaconic anhydride in refluxing toluene to provide 4-(p-methoxybenzyl)itaconate.

15

Part B:

To a solution of 3.30g (13.2 mmol) of the product from Part A in 17 mL of toluene, was added 2.08g (13.7 mmol) of 1,8-diazabicyclo[5.4.0]undec-7-ene and then 2.35g (13.7 mmol) of benzyl bromide. After 2 hours, the solution was filtered and the filtrate washed with sat'd NaHCO₃, 3N HCl, brine, dried and concentrated to afford 3.12g of an oil. After chromatography on silica gel using 0-5% ethyl acetate/hexane one obtains 2.19g (49%) of benzyl 4-(4-methoxybenzyl)itaconate.

20

25

Part C:

A solution of 1.22g (3.6 mmol) of product from Part B and 150 mg of [(R,R-Dipamp)] cyclooctadienylrhodium] tetrafluoroborate in 15 mL of methanol was thoroughly degassed, flushed with nitrogen and then hydrogen and hydrogenated under 50 psig for 16 hours. The solution was filtered and concentrated to afford 1.2g of a brown oil. This was dissolved in 5 mL of methylene chloride and 5 mL of toluene and 3 mL of trifluoroacetic

30

35

acid was added. After 4 hours, the solvents were removed in vacuo, the residue dissolved in methylene chloride, which was then extracted with sat'd NaHCO₃. After separation, the aqueous layer was acidified, reextracted with methylene chloride which was then dried and concentrated to afford 470 mg (60%) of 3(R)-methyl-4-benzylsuccinic acid.

Part D:

To a solution of 470 mg (2.11 mmol) of product from Part C and 463 mg (3.03 mg) of HOBT in 5 mL of DMF at 0°C was added 451 mg (2.35 mmol) of EDC. After 30 min. at 0°C, a solution of 728 mg (2.08 mmol) of the product from Example 34, Part E in 3 mL of DMF was added. After stirring at 0°C for 1 hour and 15 hours at RT, ethyl acetate was added and extracted with sat'd NaHCO₃, 5% aqueous citric acid, brine, dried and concentrated to give 930 mg of crude product. chromatography on silica gel using 0-10% ethyl acetate/hexane one obtains 570 mg (50%) of butanoic acid, 4-[[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-2-methyl-4-oxo-, [1S-[1R*(2S*), 2S*]-, benzyl ester.

Part E:

The product was hydrogenated in methanol using 10% palladium on carbon under 40 psig of hydrogen to afford butanoic acid, 4-[[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-2-methyl-4-oxo-, [1S-[1R*(2S*), 2S*]-.

Part F:

To a solution of 427 mg (0.92 mmol) of product from Part E and 210 mg (1.37 mmol) in 3 mL of DMF at 0°C was added 196 mg (1.02 mmol) of EDC. After 30 min. at 0°C, 2 mL of conc. aqueous ammonia was added. After 1 hour at 0°C and 15 hours at RT, ethyl acetate was added and then

extracted with sat'd NaHCO₃, brine, dried and concentrated to afford crude product. Recrystallization from ethyl acetate/hexane afforded 50 mg (12%) of butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3-methyl-, [1S-[1R*(3S*), 2S*]]-.

Example 41

10 Preparation of butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3-methyl-, [1S-[1R*(3R*), 2S*]]-.

15 This was prepared in an identical manner to the previous example except that the asymmetric hydrogenation step was done in the presence of [(S,S-dipamp)cyclooctadienyl]rhodium]-tetrafluoroborate as catalyst.

Example 42

20 Preparation of butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-, [1S-[1R*(2S*, 3R*), 2S*]]], and [1S-[1R*(2R*, 3S*), 2S*]]].

Part A:

To a solution of 863 mg (5.91 mmol) of meso-2,3-dimethylsuccinic acid in 7 mL of DMF at RT was added 1.13g (5.91 mmol) of EDC. After 15 minutes, a solution of 2.07g (5.91 mmol) of the product from Example 34, Part E and 1.4 mL of pyridine in 7 mL of anhydrous methylene chloride was added. After 11 hours, ethyl acetate was added and washed with 0.2N HCl, brine, dried and concentrated to afford 35 2.73g (97%) of a 1:1 mixture of diastereomeric acids.

Part B:

To a solution of 1.45g (3.04 mmol) of the 1:1 mixture from Part A and 613 mg (4.51 mmol) of HOBT in 10 mL of DMF at 0°C was added 635 mg (3.31 mmol) of EDC. After 30 minutes at 0°C, 5 mL of conc. aqueous ammonia was added. After 1 hour at 0°C and 14 hours at RT, ethyl acetate was added, washed with 0.2N HCl, sat'd NaHCO₃, brine, dried and concentrated to afford 0.64g (44%) of a 1:1 mixture of amides.

These were separated on a Whatman 10 micron partisil column using 8%-14% isopropanol/-methylene chloride. The first isomer to elute was identified as butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-, [1S-[1R*(2R*, 3S*), 2S*], m/e/ = 477 (M + H).

The second isomer to elute was identified as butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]-carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-, [1S-[1R*(2S*, 3R*), 2S*], m/e = 477 (M + H).

Example 43

Preparation of pentanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*, 2S*].

Part A:

To a solution of 232 mg (0.66 mmol) of the product from Example 34, Part E and 98 mg (1.2 mmol) of pyridine in 2 mL of methylene chloride was added 95 mg (0.66 mmol) of 3,3-dimethylglutaric anhydride at RT. After 15 hours, ethyl acetate was added, washed with 1N HCl, brine, dried and concentrated to afford 261 mg of crude

product. Chromatography on silica gel using 5-20% methanol/methylene chloride afforded 108 mg of acid, $m/e = 492$ ($M + H$).

5 Part B:

To a solution of 92 mg (0.19 mmol) of product from Part A and 38 mg (0.28 mmol) HOBT in 0.5 mL DMF at 0°C was added 36 mg (0.19 mmol) of EDC. After 30 minutes at 0°C, 0.25 mL of conc. aqueous ammonia was added. After 10 hour at 0°C and 16 hours at RT, ethyl acetate was added, washed with 0.2N HCl, sat'd NaHCO₃, brine, dried and concentrated to afford 72 mg of crude product. This was passed through a one-inch column of basic alumina with 10% methanol/methylene chloride to afford 53 mg of desired 15 product, $m/e = 491$ ($M + H$).

Example 44

Preparation of butanediamide, N-[3-[[[(1,1-dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-2-hydroxy-1-(phenylmethyl)propyl]-2,3-dimethyl-1S-1R*(2R*, 3S*), 2S*]] (Isomer #1) and

Preparation of butanediamide, N-[3-[[[(1,1-dimethylethyl)aminolcarbonyl](3-methylbutyl)aminol-2-hydroxy-1-(phenylmethyl)propyl]-2,3-dimethyl-1S-1R*(2R*, 3S*), 2S*]] (Isomer #2).

Part A:

30 To a solution of 1.47g (4.20 mmol) of the product from Example 34, Part E and 1.4 mL of pyridine in 9 mL of methylene chloride at RT was added 538 mg (4.20 mmol) of 2,2-dimethylsuccinic anhydride. After 15 hours, ethyl acetate was added and washed with 0.2N HCl, brine, dried 35 and concentrated to afford 1.87g of crude product (approx. 3:1 mixture of isomers).

Part B:

To a solution of 1.85g (3.9 mmol) of crude product from Part A and 887 mg (5.8 mmol) of HOBT in 10 mL of DMF at 0°C was added 809 mg (4.2 mmol) EDC. After 30 minutes at 0°C, 6 mL of conc. aqueous ammonia was added. After 1 hour at 0°C and 15 hours at RT, ethyl acetate was added, washed with 0.2N HCl, sat'd NaHCO₃, brine, dried and concentrated to afford 923 mg of crude product. The two isomers were separated on a Whatman Partisil 5 column using 8-14% isopropanol/methylene chloride. The major isomer was identified as Isomer #1, m/e = 477 (M + H).

The minor isomer was identified as Isomer #2, m/e = 477 (M + H).

15

Example 45

This example illustrates the procedure utilized to prepare compounds wherein the stereochemistry about the hydroxyl group is (S).

20

Part A:

A solution of 3(S)-(1,1-dimethylethoxycarbonyl)-amino-1,2-(R)-epoxy-4-phenylbutane (1.00g, 3.80 mmol) and isobutylamine (5.55g, 76 mmol, 20 equiv.) in 10 mL of isopropyl alcohol was warmed to 60°C for 1 hour. The solution was cooled to room temperature and concentrated in vacuo and the residue recrystallized from hexane/methylene chloride to give 0.93g, 73% of [2(S), 3(S)]-N-[[[3-[(1,1-dimethylethyl) carbamoyl]amino]]-2-hydroxy-4-phenylbutyl]N-[(3-methylbutyl)]amine, mp 91.3 - 93.0°C.

30

Part B:

The product from Part A (46.3mg, 0.14 mmol) was dissolved in a mixture of 5 mL of tetrahydrofuran and 2 mL of methylene chloride and treated with tert-butylisocyanate (136.4mg, 1.376 mmol) via syringe. The solution was stirred at room temperature for 0.5 hour and then the

35

solvent was removed in vacuo. The product, TLC on SiO₂, 1:1 hexane: ethyl acetate had R_f = 0.74 and was used directly in the next step without further purification.

5 Part C:

The crude product from Part B was taken up in 10 mL of 4N hydrochloric acid in dioxane and stirred at room temperature for 0.25 hours. The solvent and excess hydrochloric acid was removed in vacuo whereupon the product crystallized. The solid was isolated by filtration washed with acetone and dried in vacuo to 3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2(S)-hydroxy-1(S)-(phenylmethyl)propylamine hydrochloride.

15 Part D:

A solution of N-Cbz-L-asparagine (225.5mg, 0.847 mmol) and N-hydroxybenzotriazole (182.9mg, 1.21 mmol) was dissolved in 2 mL of dimethylformamide and cooled to 0°C and then treated with EDC (170.2mg, 0.898 mmol) for 10 minutes. This mixture was then treated with 3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2(S)-hydroxy-1(S)-(phenylmethyl)propylamine hydrochloride. (300.0mg, 0.807 mmol) followed by N-methylmorpholine (90.0mg, 0.888 mmol) via syringe. The solution was stirred at room temperature for 16 hours and then poured into 20 mL of rapidly stirring 60% saturated aqueous sodium bicarbonate solution whereupon a white precipitate formed. The solid was isolated by filtration, washed with saturated aqueous sodium bicarbonate solution, water, 5% aqueous citric acid solution, water and then dried in vacuo to give 319mg, 68% of butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2(S)-hydroxy-1(S)-(phenylmethyl)propyl]-2(S)-[(benzyloxycarbonyl)amino] mp 139-141°C, MH⁺ m/z = 584.

Example 46

The compounds of the present invention are effective HIV protease inhibitors. Utilizing an enzyme assay as described below, the compounds set forth in the examples herein disclosed inhibited the HIV enzyme. The preferred compounds of the present invention and their calculated IC₅₀ (inhibiting concentration 50%, i.e., the concentration at which the inhibitor compound reduces enzyme activity by 50%) values are shown in Table 16. The enzyme method is described below. The substrate is 2-aminobenzoyl-Ile-Nle-Phe(p-NO₂)-Gln-ArgNH₂. The positive control is MVT-101 (Miller, M. et al, Science, 246, 1149 (1989)). The assay conditions are as follows:

Assay buffer: 20 mM sodium phosphate, pH 6.4
20% glycerol
1 mM EDTA
1 mM DTT
0.1% CHAPS

The above described substrate is dissolved in DMSO, then diluted 10 fold in assay buffer. Final substrate concentration in the assay is 80 µM.

HIV protease is diluted in the assay buffer to a final enzyme concentration of 12.3 nanomolar, based on a molecular weight of 10,780.

The final concentration of DMSO is 14% and the final concentration of glycerol is 18%. The test compound is dissolved in DMSO and diluted in DMSO to 10x the test concentration; 10 µl of the enzyme preparation is added, the materials mixed and then the mixture is incubated at ambient temperature for 15 minutes. The enzyme reaction is initiated by the addition of 40 µl of substrate. The

increase in fluorescence is monitored at 4 time points (0, 8, 16 and 24 minutes) at ambient temperature. Each assay is carried out in duplicate wells.

5

TABLE 16

10	Compound	IC ₅₀ (nanomolar)
15	1a) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(N-benzoyloxycarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-	35
20	1b) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-	7.1
30	2a) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(N-benzoyloxycarbonyl)amino], [1S-[1R*(R*), 2S*]]-	13.4
35	2b) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-	3.2

TABLE 16 (Cont'd.)

5	Compound	IC ₅₀ (nanomolar)
10	3a) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-naphthylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(N-benzyl-oxy)carbonyl]amino]-, [1S-[1R*(R*), 2S*]]-	31.0
15	3b) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-naphthylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-	2.9
25	4a) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-phenylethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(N-benzyloxycarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-	70
30	4b) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-phenylethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-	10

TABLE 16 (Cont'd.)

5	Compound	IC ₅₀ (nanomolar)
10	5a) Butanedi am ide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](3,3-dimethylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(N-benzyl oxy carbonyl)amio], [1S-[1R*(R*),2S*]]-	18
15	5b) Butanedi am ide, N ¹ -[3-[[[(1,1-dimethylethyl)amino]carbonyl](3,3-dimethylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-	3.7
25	6) Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*),2S*]]-	29
30	7) Propanamide, 3-(acetylamino)-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-	100
35	8) Propanamide, 3-(4-methoxybenzyloxy-carbonyl)amino-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-	6

TABLE 16 (Cont'd.)

5	Compound	IC ₅₀ (nanomolar)
10	9) Butanediamide, N ¹ -[2-hydroxy-3-[(4-fluorophenylmethyl) (1-pyrrolidinyl-carbonyl) amino]-1-(phenylmethyl) propyl]-2-[(2-quinolinylcarbonyl) amino]-, [1S-[1R*(R*), 2S*]]-	7
15	10) Butanediamide, N ¹ -[3-[[[(1,1-dimethylethyl) amino] carbonyl] (3-methylbutyl) amino]-2-hydroxy-1-(phenylmethyl) propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*), 2S*]]-	12
20	11) Butanoic acid, 4-[[3-[[[(1,1-dimethylethyl) amino] carbonyl] (3-methylbutyl) amino]-2-hydroxy-1-(phenylmethyl) propyl] amino]-2,2,3-trimethyl-4-oxo-, [1S-[1R*(3S*), 2S*]]-	8
25	12) Butaneamide, 2-(acetyl amino)-N-[3-[[[(1,1-dimethylethyl) amino] carbonyl] (3-methylbutyl) amino]-2-hydroxy-1-(phenylmethyl) propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]-	18
30	13) Butaneamide, 2-[92,2-dimethylaminoacetyl) amino]-N-[3-[[[(1,1-dimethylethyl) amino] carbonyl] (3-methylbutyl) amino]-2-hydroxy-1-(phenylmethyl) propyl]-3,3dimethyl-, [1S-[1R*(R*), 2S*]]-	10
35		

Example 47

The effectiveness of the compounds listed in Table 9 were determined in the above-described enzyme assay and in a CEM cell assay.

The HIV inhibition assay method of acutely infected cells is an automated tetrazolium based colorimetric assay essentially that reported by Pawles et al, J. Virol. Methods 20, 309-321 (1988). Assays were performed in 96-well tissue culture plates. CEM cells, a CD4+ cell line, were grown in RPMI-1640 medium (Gibco) supplemented with a 10% fetal calf serum and were then treated with polybrene (2µg/ml). An 80 µl volume of medium containing 1×10^4 cells was dispensed into each well of the tissue culture plate. To each well was added a 100µl volume of test compound dissolved in tissue culture medium (or medium without test compound as a control) to achieve the desired final concentration and the cells were incubated at 37°C for 1 hour. A frozen culture of HIV-1 was diluted in culture medium to a concentration of 5×10^4 TCID₅₀ per ml (TCID₅₀ = the dose of virus that infects 50% of cells in tissue culture), and a 20µL volume of the virus sample (containing 1000 TCID₅₀ of virus) was added to wells containing test compound and to wells containing only medium (infected control cells). Several wells received culture medium without virus (uninfected control cells). Likewise, the intrinsic toxicity of the test compound was determined by adding medium without virus to several wells containing test compound. In summary, the tissue culture plates contained the following experiments:

	Cells	Drug	Virus
5			
	1. +	-	-
	2. +	+	-
	3. +	-	+
10	4. +	+	+

In experiments 2 and 4 the final concentrations of test compounds were 1, 10, 100 and 500 µg/ml. Either azidothymidine (AZT) or dideoxyinosine (ddI) was included as a positive drug control. Test compounds were dissolved in DMSO and diluted into tissue culture medium so that the final DMSO concentration did not exceed 1.5% in any case. DMSO was added to all control wells at an appropriate concentration.

Following the addition of virus, cells were incubated at 37°C in a humidified, 5% CO₂ atmosphere for 7 days. Test compounds could be added on days 0, 2 and 5 if desired. On day 7, post-infection, the cells in each well were resuspended and a 100µl sample of each cell suspension was removed for assay. A 20µL volume of a 5 mg/ml solution of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) was added to each 100µL cell suspension, and the cells were incubated for 4 hours at 27°C in a 5% CO₂ environment. During this incubation, MTT is metabolically reduced by living cells resulting in the production in the cell of a colored formazan product. To each sample was added 100µl of 10% sodium dodecylsulfate in 0.01 N HCl to lyse the cells, and samples were incubated overnight. The absorbance at 590 nm was determined for each sample using a Molecular Devices microplate reader. Absorbance values for each set of wells is compared to assess viral control infection, uninfected control cell response as well as test compound by cytotoxicity and antiviral efficacy.

TABLE 17

5	Compound	IC ₅₀	EC ₅₀	TD ₅₀
10	butanediamide N ¹ [3- [[[n-butylamino]carbonyl](2- methylpropyl)amino]-2-hydroxy- 1-(phenylmethyl)propyl]-2- [(phenylmethylcarbamoyl) amino], [1S-[1R*(R*), 2S*]]--.	940nM	17μM	170- 180μM
15	butanediamide N ¹ [3- [[[n-butylamino]carbonyl](2- methylpropyl)amino]-2-hydroxy- 1-(phenylmethyl)propyl]-2- [(2-quinolinylcarbonyl)amino]- [1S-[1R*(R*), 2S*]]-.	126nM	26nM	53μM
20	butanediamide N ¹ [3- [[[(1,1-dimethylethyl)amino]- carbonyl](2-phenylethyl)amino]- 2-hydroxy-1-(phenylmethyl)propyl]-			
25	2-[(2-quinolinylcarbonyl)amino] [1S-[1R*(R*), 2S*]]-.	10nM	28nM	13μM
30	butanediamide N ¹ [3- [[[(1,1-dimethylethyl)amino]- carbonyl](2-methylpropyl)amino]- 2-hydroxy-1-(phenylmethyl)propyl]- 2-[(2-quinolinylcarbonyl)amino] [1S-[1R*(R*), 2S*]]-.	7nM	10nM	6μM

TABLE 17 (Cont'd.)

5	Compound	IC ₅₀	EC ₅₀	TD ₅₀
10	butaneamide, 2-[92,2-dimethylaminoacetyl)amino]-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*),2S*]]-.	10nM	58nM	tba
15				

Utilizing the procedures set forth above in the examples along with the general description, it is contemplated that the compounds listed below and those shown in Tables 18 and 19 could be prepared and that such compounds would have activities as HIV protease inhibitors substantially similar to the activities of the compounds set forth in the examples.

25 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl]pentylamino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-

30 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl]hexylamino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-

35 Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl]butylamino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-fluorophenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-methoxyphenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](cyclohexylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-

Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(cyclohexylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-

Butanediamide, N¹-[2-hydroxy-3-[(3-methylbutyl)(1-pyrrolidinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-

Butanediamide, N¹-[2-hydroxy-3-[(3-methylbutyl)(1-pyrrolidinylcarbonyl)amino]-1-(4-fluorophenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-

Butanediamide, N¹-[2-hydroxy-3-[(3-methylbutyl)(1-pyrrolidinylcarbonyl)amino]-1-(cyclohexylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*), 2S*]]-

Butanediamide, N¹-[2-hydroxy-3-[(phenylmethyl)(4-morpholinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-

- 5 Butanediamide, N¹-[2-hydroxy-3-[(3-methylbutyl)(1-piperidinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-

- 10 Butanediamide, N¹-[2-hydroxy-3-[(phenylmethyl)(1-piperidinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-

- 15 Butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl]butylamino]-2-hydroxy-1-(phenylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-

- 20 Butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-

- 25 Butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-fluorophenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-

- 30 Butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](cyclohexylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-

- 35 Butanediamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(cyclohexylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-

Butanoic acid, 4-[[3-[[[(1,1-
dimethylethyl)amino]carbonyl]butylamino]-2-hydroxy-1-
(phenylmethyl)propyl]amino-2,2,3-trimethyl-4-oxo-, [1S-
[1R*(3S*),2S*]]-

5

Butanoic acid, 4-[[3-[[[(1,1-
dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl]amino-2,2,3-trimethyl-4-
oxo-, [1S-[1R*(3S*),2S*]]-

10

Butanoic acid, 4-[[3-[[[(1,1-
dimethylethyl)amino]carbonyl](4-fluorophenylmethyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl]amino-2,2,3-trimethyl-4-
oxo-, [1S-[1R*(3S*),2S*]]-

15

Butanoic acid, 4-[[3-[[[(1,1-
dimethylethyl)amino]carbonyl](cyclohexylmethyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl]amino-2,2,3-trimethyl-4-
oxo-, [1S-[1R*(3S*),2S*]]-

20

Butaneamide,
2-(acetylamino)-N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
hydroxy-1-(cyclohexylmethyl)propyl]-3,3-dimethyl-, [1S-
[1R*(R*),2S*]]-

25

Butaneamide,
2-(acetylamino)-N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl]butylamino]-2-hydroxy-1-
(phenylmethyl)propyl]-3,3-dimethyl-, [1S-[1R*(R*),2S*]]-

30

Butaneamide,
2-(acetylamino)-N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-
[1R*(R*),2S*]]-

35

Butaneamide,

2-(acetyl amino)-N-[3-[[[(1,1-
dimethylethyl) amino] carbonyl] (cyclohexylmethyl) amino]-2-
hydroxy-1-(phenylmethyl) propyl]-3,3-dimethyl-, [1S-
5 [1R*(R*), 2S*]]-

Butaneamide,

2-[(2,2-dimethylaminoacetyl) amino]-N-[3-[[[(1,1-
dimethylethyl) amino] carbonyl] (3-methylbutyl) amino]-2-
10 hydroxy-1-(phenylmethyl) propyl]-3-methyl-, [1S-
[1R*(R*), 2S*]]-

Butaneamide,

2-[(2,2-dimethylaminoacetyl) amino]-N-[3-[[[(1,1-
15 dimethylethyl) amino] carbonyl] (3-methylbutyl) amino]-2-
hydroxy-1-(cyclohexylmethyl) propyl]-3,3-dimethyl-, [1S-
[1R*(R*), 2S*]]-

Butaneamide,

20 2-[(2,2-dimethylaminoacetyl) amino]-N-[3-[[[(1,1-
dimethylethyl) amino] carbonyl] butylamino]-2-hydroxy-1-
(phenylmethyl) propyl]-3,3-dimethyl-, [1S-[1R*(R*), 2S*]]-

Butaneamide,

25 2-[(2,2-dimethylaminoacetyl) amino]-N-[3-[[[(1,1-
dimethylethyl) amino] carbonyl] (phenylmethyl) amino]-2-
hydroxy-1-(phenylmethyl) propyl]-3,3-dimethyl-, [1S-
[1R*(R*), 2S*]]-

30 Butaneamide,


2-[(2,2-dimethylaminoacetyl) amino]-N-[3-[[[(1,1-
dimethylethyl) amino] carbonyl] (4-fluorophenylmethyl) amino]-
2-hydroxy-1-(phenylmethyl) propyl]-3,3-dimethyl-, [1S-
[1R*(R*), 2S*]]-

35

Butaneamide,
 2-[(2,2-dimethylaminoacetyl)amino]-N-[3-[[[(1,1-
 dimethylethyl)amino]carbonyl](cyclohexylmethyl)amino]-2-
 hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-
 5 [1R*(R*),2S*]]-

TABLE 18

10



15

	R ¹	R ²	R ³
	t-Butyl	Benzyl	p-Fluorobenzyl
	i-Butyl	Benzyl	i-Amyl
	i-Propyl	Benzyl	i-Amyl
	Propargyl	Benzyl	i-Amyl
20	t-Butyl	Benzyl	i-Amyl
	t-Butyl	Benzyl	Benzyl
	t-Butyl	Benzyl	n-Butyl
	sec-Butyl	Benzyl	i-Amyl
	C(CH ₃) ₂ (SCH ₃)	Benzyl	i-Amyl
25	t-Butyl	p-Fluorobenzyl	p-Methoxybenzyl
	i-Butyl	p-Fluorobenzyl	i-Amyl
	i-Propyl	p-Fluorobenzyl	i-Amyl
	Propargyl	p-Fluorobenzyl	i-Amyl
	t-Butyl	p-Fluorobenzyl	i-Amyl
30	t-Butyl	p-Fluorobenzyl	Benzyl
	t-Butyl	p-Fluorobenzyl	n-Butyl
	sec-Butyl	Benzyl	i-Amyl
	C(CH ₃) ₂ (SCH ₃)	Benzyl	i-Amyl
	t-Butyl	Cyclohexylmethyl	p-Fluorobenzyl
35	i-Butyl	Cyclohexylmethyl	i-Amyl
	i-Propyl	Cyclohexylmethyl	i-Amyl
	Propargyl	Cyclohexylmethyl	i-Amyl

TABLE 18 (Cont'd.)

5	R ¹	R ²	R ³
	t-Butyl	Cyclohexylmethyl	i-Amyl
	t-Butyl	Cyclohexylmethyl	Benzyl
10	t-Butyl	Cyclohexylmethyl	n-Butyl
	sec-Butyl	Cyclohexylmethyl	i-Amyl
	C(CH ₃) ₂ (SCH ₃)	Cyclohexylmethyl	i-Amyl
	t-Butyl	n-Butyl	Cyclohexylmethyl
	i-Butyl	n-Butyl	i-Amyl
15	i-Propyl	n-Butyl	i-Amyl
	Propargyl	n-Butyl	i-Amyl
	t-Butyl	n-Butyl	i-Amyl
	t-Butyl	n-Butyl	Benzyl
	t-Butyl	n-Butyl	n-Butyl
20	sec-Butyl	n-Butyl	i-Amyl
	C(CH ₃) ₂ (SCH ₃)	n-Butyl	i-Amyl

TABLE 19

5			
	R^1	R^2	R^3
10	methyl benzyl hydroxymethyl 1-hydroxyethyl	methyl methyl methyl methyl	methyl methyl methyl methyl
15	hydrogen hydrogen hydrogen	methyl methyl $R^2 + R^3 = \text{imidazole}$	ethyl benzyl

20 The compounds of the present invention are effective antiviral compounds and, in particular, are effective inhibitors of retroviruses, particularly,

25 lentoviruses as shown above. Thus, the subject compounds are effective inhibitors of HIV. It is contemplated that the subject compounds will also inhibit other strains of

30 HIV, such as HIV-2 and other viruses such as, for example, human T-cell leukemia virus, feline leukemia virus, simian immunodeficiency virus, respiratory syncytial virus,

35 hepadnavirus, cytomegalovirus and picornavirus. Thus, the subject compounds are effective in the treatment and/or prophylaxis of retroviral infections.

35 Compounds of the present can possess one or more asymmetric carbon atoms and are thus capable of existing in the form of optical isomers as well as in the form of

40 racemic or nonracemic mixtures thereof. The optical

isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example by formation of diastereoisomeric salts by treatment with an optically active acid or base. Examples of appropriate acids are tartaric, diacetyltartaric, dibenzoyltartaric, ditoluoyltartaric and camphorsulfonic acid and then separation of the mixture of diastereoisomers by crystallization followed by liberation of the optically active bases from these salts. A different process for separation of optical isomers involves the use of a chiral chromatography column optimally chosen to maximize the separation of the enantiomers. Still another available method involves synthesis of covalent diastereoisomeric molecules by reacting compounds of Formula I with an optically pure acid in an activated form or an optically pure isocyanate. The synthesized diastereoisomers can be separated by conventional means such as chromatography, distillation, crystallization or sublimation, and then hydrolyzed to deliver the enantiomerically pure compound. The optically active compounds of Formula I can likewise be obtained by utilizing optically active starting materials. These isomers may be in the form of a free acid, a free base, an ester or a salt.

The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, mesylate and undecanoate.

Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, 5 and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

10

Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, 15 maleic acid, succinic acid and citric acid. Other examples include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic bases.

Total daily dose administered to a host in single or divided doses may be in amounts, for example, from 0.001 to 10 mg/kg body weight daily and more usually 0.01 to 1 mg. Dosage unit compositions may contain such amounts of submultiples thereof to make up the daily dose.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

The dosage regimen for treating a disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex, diet and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetic and toxicology profiles of the particular compound employed, whether a drug delivery system is utilized and whether the compound is administered as part of a drug combination. Thus, the dosage regimen actually employed may vary widely and therefore may deviate from the preferred dosage regimen set forth above.

25

The compounds of the present invention may be administered orally, parenterally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection, or infusion techniques.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Suppositories for rectal administration of the drug can be prepared by mixing the drug with a suitable nonirritating excipient such as cocoa butter and polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum and release the drug.

Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration may include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions
5 may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

While the compounds of the invention can be
10 administered as the sole active pharmaceutical agent, they can also be used in combination with one or more immunomodulators, antiviral agents or other antiinfective agents. For example, the compounds of the invention can be administered in combination with AZT or with N-butyl-1-
15 deoxynojirimycin for the prophylaxis and/or treatment of AIDS. When administered as a combination, the therapeutic agents can be formulated as separate compositions which are given at the same time or different times, or the therapeutic agents can be given as a single composition.


20

The foregoing is merely illustrative of the invention and is not intended to limit the invention to the disclosed compounds. Variations and changes which are obvious to one skilled in the art are intended to be within
25 the scope and nature of the invention which are defined in the appended claims.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics
30 of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.


WHAT IS CLAIMED IS:

1. A compound represented by the formula:




5


wherein A represents radicals represented by formulas:



(A1)



(A2)



(A3)

10

wherein R represents hydrogen, alkoxycarbonyl, aralkoxycarbonyl, alkylcarbonyl, cycloalkylcarbonyl, cycloalkylalkoxycarbonyl, cycloalkylalkanoyl, alkanoyl, aralkanoyl, aroyl, aryloxy carbonyl, aryloxy carbonylalkyl, aryloxyalkanoyl, heterocyclylcarbonyl, heterocyclyloxy carbonyl, heterocyclylalkanoyl, heterocyclylalkoxycarbonyl, heteroaralkoxycarbonyl, heteroaryloxy carbonyl, heteroaroyl, heteroaralkanoyl, alkyl, aryl, aralkyl, aryloxyalkyl, heteroaryloxyalkyl, hydroxyalkyl, aminocarbonyl, aminoalkanoyl, and mono- and disubstituted aminocarbonyl and mono- and disubstituted aminoalkanoyl radicals wherein the substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl,

15

20

heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkyl radicals, or in the case of a disubstituted aminoalkanoyl radical, said substituents along with the nitrogen atom to which they are attached form a heterocyclyl or heteroaryl radical;

R' represents hydrogen and radicals as defined for R³ or R and R' together with the nitrogen to which they are attached form a heterocycloalkyl or heteroaryl radical, or when A is Al, R' represents hydrogen, radicals as defined for R³ and arakloxy carbonylalkyl and aminocarbonylalkyl and aminocarbonyl radicals wherein said amino group may be mono- or disubstituted with substituents selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl and heterocycloalkylalkyl radicals;

t represents 0 or 1;

R¹ represents hydrogen, -CH₂SO₂NH₂, -CO₂CH₃, -CONHCH₃, -CON(CH₃)₂, -CH₂C(O)NHCH₃, -CH₂C(O)N(CH₃)₂, -CONH₂, -C(CH₃)₂(SCH₃), -C(CH₃)₂(S[O]CH₃), -C(CH₃)₂(S[O]CH₃), alkyl, haloalkyl, alkenyl, alkynyl and cycloalkyl radicals and amino acid side chains selected from asparagine, S-methyl cysteine and the corresponding sulfoxide and sulfone derivatives thereof, glycine, leucine, isoleucine, allo-isoleucine, tert-leucine, phenylalanine, ornithine, alanine, histidine, norleucine, glutamine, valine, threonine, serine, aspartic acid, beta-cyano alanine, and allothreonine side chains;

R^{1'} and R^{1''} independently represent hydrogen and radicals as defined for R¹, or one of R^{1'} and R^{1''}, together with R¹ and the carbon atoms to which they are attached, represent a cycloalkyl radical;

R² represents alkyl, aryl, cycloalkyl, cycloalkylalkyl and aralkyl radicals optionally substituted with a group selected from -NO₂, -OR⁹, -SR⁹, and halogen radicals, wherein R⁹ represents hydrogen and alkyl radicals;

5

R³ represents alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, heterocycloalkylalkyl, aryl, aralkyl, heteroaralkyl, aminoalkyl and mono- and disubstituted aminoalkyl radicals, wherein said substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkyl radical, said substituents along with the nitrogen atom to which they are attached, form a heterocycloalkyl or a heteroaryl radical;

10

15

X' represents O, N and C(R¹⁷) wherein R¹⁷ represents hydrogen and alkyl radicals;

20

Y, Y' and Y" independently represent O and S;

R⁴ and R⁵ independently represent hydrogen and radicals as defined by R³, or R⁴ and R⁵ together with the nitrogen atom to which they are bonded represent heterocycloalkyl and heteroaryl radicals;

25

R⁶ represents hydrogen and alkyl radicals;

30


R²⁰, R²¹, R³⁰, R³¹ and R³² represent radicals as defined for R¹, or one of R¹ and R³⁰ together with one of R³¹ and R³² and the carbon atoms to which they are attached form a cycloalkyl radical; or R³⁰ and R³² together with the carbon atoms to which they are attached form a three to six-membered cycloalkyl radical; and

35

R³³ and R³⁴ independently represent hydrogen and radicals as defined for R³, or R³³ and R³⁴ together with X' represent cycloalkyl, aryl, heterocyclyl and heteroaryl radicals, provided that when X' is O, R³⁴ is absent.

5

2. Compound represented by the formula:



10 wherein; R represents alkoxy carbonyl, aralkoxy carbonyl, alkyl carbonyl, cycloalkyl carbonyl, cycloalkyl alkoxy carbonyl, cycloalkyl alkanoyl, alkanoyl, cycloalkanoyl, cycloalkyl alkanoyl, aralkanoyl, aroyl, aryloxy carbonyl, aryloxy carbonyl alkyl, aryloxy alkanoyl, heterocyclocarbonyl, heterocyclyloxy carbonyl, heterocyclyl, 15 alkanoyl, heterocyclyl alkoxy carbonyl, heteroaralkanoyl, heteroaralkoxy carbonyl, heteroaryloxy carbonyl, heteroaroyl, alkyl, alkenyl, cycloalkyl, aryl, aralkyl, aryloxy alkyl, heteroaryloxy alkyl, hydroxy alkyl, aminocarbonyl, and 20 aminoalkanoyl radicals, and mono- and disubstituted aminoalkanoyl and mono- and disubstituted aminocarbonyl radicals wherein the substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkyl alkyl, heteroaryl, heteroaralkyl, heteroalkyl, heterocycloalkyl alkyl radicals, 25 or where said aminoalkanoyl radical is disubstituted, said substituents along with the nitrogen atom to which they are attached, form a heterocycloalkyl or heteroaryl radical;

30 R' represents hydrogen and radicals as defined for R³, or R and R' together with the nitrogen to which they are attached form a heterocycloalkyl or heteroaryl radical;

R¹ represents hydrogen, -CH₂SO₂NH₂, -CH₂CO₂CH₃, -CO₂CH₃, 35 -CONH₂, -CH₂C(O)NHCH₃, -C(CH₃)₂(SCH₃), -C(CH₃)₂(S[O]CH₃),

- C(CH₃)₂(S[O]₂CH₃), alkyl, haloalkyl, alkenyl, alkynyl and cycloalkyl radicals, and amino acid side chains selected from asparagine, S-methyl cysteine and the sulfoxide (SO) and sulfone (SO₂) derivatives thereof, isoleucine,
- 5 allo-isoleucine, alanine, leucine, tert-leucine, phenylalanine, ornithine, histidine, norleucine, glutamine, threonine, glycine, allo-threonine, serine, aspartic acid, beta-cyano alanine and valine side chains;
- 10 R^{1'} and R^{1''} independently represent hydrogen and radicals as defined for R¹;
- R² represents alkyl, aryl, cycloalkyl, cycloalkylalkyl, and aralkyl radicals, which radicals are optionally
- 15 substituted with a group selected from halogen radicals and NO₂, -OR⁹ and SR⁹ wherein R⁹ represents hydrogen and alkyl radicals;
- R³ represents alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, heterocycloalkylalkyl, aryl, aralkyl, heteroaralkyl, aminoalkyl and mono- and disubstituted aminoalkyl radicals, wherein said substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl,
- 20 heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkyl radical, said substituents along with the nitrogen atom to which they are attached, form a heterocycloalkyl or a heteroaryl radical;
- 25 R⁴ and R⁵ independently represent hydrogen and radicals as defined by R³, or together with the nitrogen atom to which they are bonded represent a heterocycloalkyl or heteroaryl radical;
- 30 t represents 0 or 1; and
- 35

Y and Y' independently represent O and S.

3. Compound of Claim 2 wherein R represents aralkoxycarbonyl and heteroaroyl radicals.

5

4. Compound of Claim 2 wherein R represents carbobenzoxy, 2-benzofurancarbonyl and 2-quinolinylcarbonyl radicals.

10

5. Compound of Claim 2 wherein R represents carbobenzoxy.

6. Compound of Claim 2 wherein R represents 2-quinolinylcarbonyl and 2-benzofurancarbonyl radicals.

15

7. Compound of Claim 2 wherein R¹ represents alkyl, alkenyl and alkynyl radicals, and amino acid side chains selected from the group consisting of asparagine, valine, threonine, allo-threonine, isoleucine, S-methyl cysteine and the sulfone and sulfoxide derivatives thereof, alanine, and allo-isoleucine.

20

8. Compound of Claim 2 wherein R¹ represents methyl, propargyl, t-butyl, isopropyl and sec-butyl radicals, and amino acid side chains selected from the group consisting of asparagine, valine, S-methyl cysteine, allo-iso-leucine, iso-leucine, threonine, serine, aspartic acid, beta-cyano alanine, and allo-threonine side chains.

25

9. Compound of Claim 2 wherein R¹ represents methyl and t-butyl radicals.

30

10. Compound of Claim 2 wherein R¹ represents a t-butyl radical when t is O.

35

11. Compound of Claim 2 wherein R¹ represents amino acid side chains selected from asparagine, valine, alanine and isoleucine side chains.

5 12. Compound of Claim 2 wherein R¹ represents amino acid side chains selected from asparagine, iso-leucine and valine side chains.

10 13. Compound of Claim 2 wherein R¹ represents an asparagine side chain.

14. Compound of Claim 2 wherein R¹ represents a t-butyl radical and an asparagine side chain.

15 15. Compound of Claim 2 wherein R¹ represents a methyl radical when t is 1.

16. Compound of Claim 2 wherein t is 0.

20 17. Compound of Claim 2 wherein t is 1.

18. Compound of Claim 2 wherein R² represents alkyl, cycloalkylalkyl and aralkyl radicals, which radicals are optionally substituted with halogen radicals and
25 radicals represented by the formula -OR⁹ and -SR⁹ wherein R⁹ represents hydrogen and alkyl radicals.

19. Compound of Claim 2 wherein R² represents alkyl, cycloalkylalkyl and aralkyl radicals.

30 20. Compound of Claim 2 wherein R² represents aralkyl radicals.

21. Compound of Claim 2 wherein R² represents
35 CH₃SCH₂CH₂-, iso-butyl, n-butyl, benzyl, 4-fluorobenzyl, 2-naphthylmethyl and cyclohexylmethyl radicals.

22. Compound of Claim 2 wherein R² represents an n-butyl and iso-butyl radicals.

23. Compound of Claim 2 wherein R² represents
5 benzyl, 4-fluorobenzyl and 2-naphthylmethyl radicals.

24. Compound of Claim 2 wherein R² represents a cyclohexylmethyl radical.

10 25. Compound of Claim 2 wherein R³, R⁴ and R⁵ independently represent alkyl, alkenyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, aralkyl and heteroaralkyl radicals.

15 26. Compound of Claim 25 wherein R⁵ represents hydrogen.

20 27. Compound of Claim 25 wherein R³ and R⁴ independently represent alkyl and alkenyl radicals.

28. Compound of Claim 26 wherein R³ and R⁴ independently represent alkyl and hydroxyalkyl radicals.

25 29. Compound of Claim 26 wherein R³ and R⁴ independently represent alkyl, cycloalkyl and cycloalkylalkyl radicals.

30 30. Compound of Claim 26 wherein R³ and R⁴ independently represent alkyl, heterocycloalkyl and heterocycloalkylalkyl radicals.

35 31. Compound of Claim 26 wherein R³ and R⁴ independently represent alkyl, aryl and aralkyl radicals.

32. Compound of Claim 26 wherein R⁴ represents a t-butyl, ethyl, isopropyl and 1,1-dimethylpropyl radicals.

5 33. Compound of Claim 2 wherein R³ represents alkyl radicals having from about 2 to about 5 carbon atoms.

 34. Compound of Claim 2 wherein R³ represents n-pentyl, n-hexyl, n-propyl, i-butyl, neo-pentyl, i-amyl,
10 and n-butyl radicals.

 35. Compound of Claim 2 wherein R³, R⁴ and R⁵ independently represent alkyl radicals having from about 2 to about 5 carbon atoms, cycloalkylalkyl radicals, aralkyl
15 radicals, heterocycloalkylalkyl radicals and heteroaralkyl radicals.

 36. Compound of Claim 2 wherein R³ represents benzyl, para-fluorobenzyl, para-methoxybenzyl, ~~para-~~
20 methylbenzyl, and 2-naphthylmethyl radicals and R⁴ represents t-butyl.

 37. Compound of Claim 2 wherein R³ is cyclohexylmethyl and R⁴ is t-butyl.
25

 38. Compound of Claim 2 wherein R³ is i-amyl and R⁴ is t-butyl.

 39. Compound of Claim 2 wherein R³ is i-butyl
30 and R⁴ is t-butyl.

 40. Compound of Claim 2 wherein R³ is n-butyl and R⁴ is t-butyl.

 41. Compound of Claim 2 wherein R³ is neo-pentyl and R⁴ is t-butyl.
35

42. Compound of Claim 2 wherein R⁴ represents alkyl and cycloalkyl radicals.

5 43. Compound of Claim 2 wherein R⁵ represents hydrogen and alkyl and cycloalkyl radicals.

10 44. Compound of Claim 2 wherein R⁴ and R⁵ together with the nitrogen to which they are bonded represent pyrrolidinyl, piperidinyl, morpholinyl, and piperazinyl radicals.

45. Compound of Claim 2 wherein R³ represents heteroaralkyl radicals and R⁴ is t-butyl.

15 46. Compound of Claim 2 wherein R³ is a p-fluorobenzyl radical and R⁴ is a t-butyl radical.

20 47. Compound of Claim 2 wherein R³ is a 4-pyridylmethyl radical or its N-oxide and R⁴ is a t-butyl radical.

25 48. Compound of Claim 2 wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached represent a 5 to 8-membered heterocyclic or heterobicyclicalkyl radical.

30 49. Compound of Claim 2 wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached represent a 5 to 8-membered heterocyclyl radical, optionally substituted with an alkyl radical having from 1 to about 3 carbon atoms.

35 50. Compound of Claim 2 wherein R¹ and R^{1'} are both hydrogen and R^{1''} represents an alkyl radical having from 1 to about 4 carbon atoms.

51. Compound of Claim 2 wherein R¹ and R^{1'} are both hydrogen and R^{1"} represents -CH₂SO₂NH₂, CO₂NH₂, CO₂CH₃, alkyl, alkynyl and cycloalkyl radicals and amino acid side chains selected from asparagine, S-methyl cysteine and the sulfone and sulfoxide derivatives thereof, histidine, norleucine, glutamine, glycine, allo-isoleucine, alanine, threonine, isoleucine, leucine, tert-leucine, phenylalanine, ornithine, allo-threonine, serine, aspartic acid, beta-cyano alanine and valine side chains.
52. Compound of Claim 2 where t is O and R¹ represents the amino acid side chain of asparagine.
53. Compound of Claim 52 where R represents a heteroaroyl radical.
54. Compound of Claim 52 where R represents a 2-quinolinylcarbonyl or 2-benzofurancarbonyl radical.
55. Compound of Claim 2 where t is O and R¹ represents a t-butyl or propargyl radical or an amino acid side chain of valine or isoleucine.
56. Compound of Claim 55 where R represents an arylalkanoyl, aryloxy carbonyl, alkanoyl, aminocarbonyl, mono-substituted aminoalkanoyl, disubstituted aminoalkanoyl, or mono- or dialkylaminocarbonyl radicals.
57. Compound of Claim 55 where R represents an arylalkanoyl, aryloxy carbonyl or alkanoyl radical.
58. Compound of Claim 55 where R represents an aminocarbonyl radical, a monosubstituted aminoalkanoyl radical or disubstituted aminoalkanoyl radical.

59. Compound of Claim 55 where R represents acetyl, N,N-dimethylaminoacetyl or N-benzyl-N-methylaminoacetyl.

5 60. Compound of Claim 2 where t is 1 and R¹ is a methyl radical.

61. Compound of Claim 60 where R represents an alkanoyl, arylalkanoyl, aryloxyalkanoyl or
10 arylalkyloxy carbonyl radical.

62. Compound of Claim 60 where R represents a phenoxyacetyl, 2-naphthyloxyacetyl, benzyloxycarbonyl or p-methoxybenzyloxycarbonyl radical.

15 63. Compound of Claim 2 wherein t is 1, R¹ and R^{1'} are both hydrogen and R^{1''} is CONH₂.

64. Compound of Claim 60 where R represents an
20 aminocarbonyl or an alkylaminocarbonyl radical.

65. Compound of Claim 60 where R represents an N-methylaminocarbonyl radical.

25 66. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

67. A pharmaceutical composition comprising a
30 compound of Claim 2 and a pharmaceutically acceptable carrier.

68. Method of inhibiting a retroviral protease comprising administering a protease inhibiting amount of a
35 composition of Claim 66.

69. Method of Claim 68 wherein the retroviral protease is HIV protease.

70. Method of treating a retroviral infection comprising administering an effective amount of a composition of Claim 66.

71. Method of Claim 70 wherein the retroviral infection is an HIV infection.

72. Method for treating AIDS comprising administering an effective amount of a composition of Claim 66.

73. Method of inhibiting a retroviral protease comprising administering a protease inhibiting amount of a composition of Claim 67.


74. Method of Claim 73 wherein the retroviral protease is HIV protease.

75. Method of treating a retroviral infection comprising administering an effective amount of a composition of Claim 67.

76. Method of Claim 75 wherein the retroviral infection is an HIV infection.

77. Method for treating AIDS comprising administering an effective amount of a composition of Claim 67.

78. Compound represented by the formula:



wherein R' represents alkyl, alkenyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, aralkyl, aralkoxycarbonylalkyl, heteroaralkyl, aminoalkyl and aminocarbonylalkyl radicals, wherein the amino group of said aminoalkyl and aminocarbonylalkyl radicals may be mono- or di-substituted with substituents selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl and heteroaralkyl radicals;

R¹ represents hydrogen, -CH₂SO₂NH₂, -CO₂CH₃, -CONHCH₃, -CON(CH₃)₂, -CH₂C(O)NHCH₃, -CH₂C(O)N(CH₃)₂, -CONH₂, -C(CH₃)₂(SCH₃), -C(CH₃)₂(S[O]CH₃), -C(CH₃)₂(S[O]CH₃), alkyl, haloalkyl, alkenyl, alkynyl and cycloalkyl radicals and amino acid side chains selected from asparagine, S-methyl cysteine and the corresponding sulfoxide and sulfone derivatives thereof, glycine, leucine, isoleucine, allo-isoleucine, tert-leucine, phenylalanine, ornithine, alanine, histidine, norleucine, glutamine, valine, threonine, serine, aspartic acid, beta-cyano alanine, and allothreonine side chains;

R² represents alkyl, aryl, cycloalkyl, cycloalkylalkyl and aralkyl radicals optionally substituted with a group selected from -OR⁹, -SR⁹, and halogen radicals, wherein R⁹ represents hydrogen and alkyl radicals;

R³ represents alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, heterocycloalkylalkyl, aryl, aralkyl, heteroaralkyl, aminoalkyl and mono- and disubstituted aminoalkyl radicals, wherein said substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkyl radical, said substituents along

with the nitrogen atom to which they are attached, form a heterocycloalkyl or a heteroaryl radical;

5 R⁴ and R⁵, independently represent hydrogen and radicals as defined by R³, or R⁴ and R⁵ together with the nitrogen atom to which they are bonded represent heterocycloalkyl and heteroaryl radicals; and

10 R²⁰ and R²¹ represent radicals as defined for R¹; and Y and Y' independently represent O and S.

79. Compound of Claim 78 wherein t is O.

15 80. Compound of Claim 78 wherein R¹ represents hydrogen, alkyl and alkynyl radicals.

20 81. Compound of Claim 78 wherein R¹ represents alkyl radicals having from 1 to about 4 carbon atoms and alkenyl radicals having from 3 to 8 carbon atoms.

82. Compound of Claim 78 wherein R¹ represents methyl, ethyl, isopropyl and t-butyl radicals.

25 83. Compound of Claim 78 wherein R²⁰ and R²¹ independently represent hydrogen and alkyl radicals.

84. Compound of Claim 78 wherein R²⁰ and R²¹ independently represent hydrogen and methyl radicals.

30 85. Compound of Claim 78 wherein R²⁰ is hydrogen and R²¹ is an alkyl radical.

35 86. Compound of Claim 78 wherein R' represents alkyl, aryl and aralkyl radicals.

87. Compound of Claim 78 wherein R' is selected from methyl and phenethyl radicals.

5 88. Compound of Claim 78 wherein R² represents alkyl, cycloalkylalkyl and aralkyl radicals, which radicals are optionally substituted with halogen radicals and radicals represented by the formula -OR⁹ and -SR⁹ wherein R⁹ represents hydrogen and alkyl radicals.

10 89. Compound of Claim 78 wherein R² represents alkyl, cycloalkylalkyl and aralkyl radicals.

15 90. Compound of Claim 78 wherein R² represents aralkyl radicals.

91. Compound of Claim 78 wherein R² represents CH₃SCH₂CH₂-, iso-butyl, n-butyl, benzyl, 4-fluorobenzyl, 2-naphthylmethyl and cyclohexylmethyl radicals.

20 92. Compound of Claim 78 wherein R² represents an n-butyl and iso-butyl radicals.

25 93. Compound of Claim 78 wherein R² represents benzyl, 4-fluorobenzyl, and 2-naphthylmethyl radicals.

94. Compound of Claim 78 wherein R² represents a cyclohexylmethyl radical.

30 95. Compound of Claim 78 wherein R³, R⁴ and R⁵ independently represent alkyl, alkenyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, aralkyl and heteroaralkyl radicals.

35 96. Compound of Claim 95 wherein R⁵ represents hydrogen.

97. Compound of Claim 95 wherein R³ and R⁴ independently represent alkyl and alkenyl radicals.

5 98. Compound of Claim 96 wherein R³ and R⁴ independently represent alkyl and hydroxyalkyl radicals.

99. Compound of Claim 96 wherein R³ and R⁴ independently represent alkyl, cycloalkyl and cycloalkylalkyl radicals.

10

100. Compound of Claim 96 wherein R³ and R⁴ independently represent alkyl, heterocycloalkyl and heterocycloalkylalkyl radicals.

15

101. Compound of Claim 96 wherein R³ and R⁴ independently represent alkyl, aryl and aralkyl radicals.

102. Compound of Claim 96 wherein R³ and R⁴ independently represent alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, aralkyl, and heteroaralkyl radicals.

20

103. Compound of Claim 78 wherein R³ represents alkyl radicals having from about 2 to about 5 carbon atoms.

25

104. Compound of Claim 96 wherein R⁴ represents ethyl, i-propyl, t-butyl and 1,1-dimethylpropyl radicals.

105. Compound of Claim 78 wherein R³, R⁴ and R⁵ independently represent alkyl radicals having from about 2 to about 5 carbon atoms, cycloalkylalkyl radicals, aralkyl radicals, heterocycloalkylalkyl radicals and heteroaralkyl radicals.

30

106. Compound of Claim 78 wherein R³ represents benzyl, para-fluorobenzyl, para-methoxybenzyl, para-methylbenzyl, and 2-naphthylmethyl radicals and R⁴ represents t-butyl.

5

107. Compound of Claim 78 wherein R³ is cyclohexylmethyl and R⁴ is t-butyl.

108. Compound of Claim 78 wherein R³ is i-amyl and R⁴ is t-butyl.

10

109. Compound of Claim 78 wherein R³ is i-butyl and R⁴ is t-butyl.

15

110. Compound of Claim 78 wherein R³ is n-butyl and R⁴ is t-butyl.

111. Compound of Claim 78 wherein R³ is neopentyl and R⁴ is t-butyl.

20

112. Compound of Claim 78 wherein R⁴ represents alkyl and cycloalkyl radicals.

113. Compound of Claim 78 wherein R⁵ represents hydrogen and alkyl and cycloalkyl radicals.

25

114. Compound of Claim 78 wherein R⁴ and R⁵ together with the nitrogen to which they are bonded represent pyrrolidinyl, piperidinyl, morpholinyl, and piperazinyl radicals.

30

115. Compound of Claim 78 wherein R³ represents heteroaralkyl radicals and R⁴ is t-butyl.

116. Compound of Claim 78 wherein R³ is selected from p-fluorobenzyl, benzyl and p-methoxybenzyl radicals and R⁴ is a t-butyl radical.

35

117. A pharmaceutical composition comprising a compound of Claim 78 and a pharmaceutically acceptable carrier.

5

118. Method of inhibiting a retroviral protease comprising administering a protease inhibiting amount of a composition of Claim 117.

10

119. Method of Claim 118 wherein the retroviral protease is HIV protease.

15

120. Method of treating a retroviral infection comprising administering an effective amount of a composition of Claim 117.


121. Method of Claim 120 wherein the retroviral infection is an HIV infection.

20

122. Method for treating AIDS comprising administering an effective amount of a composition of Claim 117.

25

123. Compound represented by the formula:



wherein: R^1 represents hydrogen, $-CH_2SO_2NH_2$, $-CO_2CH_3$, $-CONHCH_3$, $-CON(CH_3)_2$, $-CH_2C(O)NHCH_3$, $-CH_2C(O)N(CH_3)_2$, $-CONH_2$, $-C(CH_3)_2(SCH_3)$, $-C(CH_3)_2(S[O]CH_3)$, $-C(CH_3)_2(S[O]CH_3)$, alkyl, haloalkyl, alkenyl, alkynyl and cycloalkyl radicals and amino acid side chains selected from asparagine, S-methyl cysteine and the corresponding sulfoxide and sulfone derivatives thereof, glycine, leucine, isoleucine,

30

allo-isoleucine, tert-leucine, phenylalanine, ornithine, alanine, histidine, norleucine, glutamine, valine, threonine, serine, aspartic acid, beta-cyano alanine, and allothreonine side chains;

5

R² represents alkyl, aryl, cycloalkyl, cycloalkylalkyl and aralkyl radicals optionally substituted with a group selected from -OR⁹, -SR⁹, and halogen radicals, wherein R⁹ represents hydrogen and alkyl radicals;

10

R³ represents alkyl, alkenyl, alkynyl, hydroxyalkyl, alkoxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heteroaryl, heterocycloalkylalkyl, aryl, aralkyl, heteroaralkyl, aminoalkyl and mono- and disubstituted aminoalkyl radicals, wherein said substituents are selected from alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocycloalkyl, and heterocycloalkylalkyl radicals, or in the case of a disubstituted aminoalkyl radical, said substituents along with the nitrogen atom to which they are attached, form a heterocycloalkyl or a heteroaryl radical;

20

R⁴ and R⁵, independently represent hydrogen and radicals as defined by R³, or R⁴ and R⁵, together with the nitrogen atom to which they are bonded represent heterocycloalkyl and heteroaryl radicals;

25

R³⁰, R³¹ and R³² represent radicals as defined for R¹, or one of R¹ and R³⁰ together with one of R³¹ and R³² and the carbon atoms to which they are attached form a cycloalkyl radical; or R³⁰ and R³² together with the carbon atoms to which they are attached form a three to six-membered cycloalkyl radical;

30

R³³ and R³⁴ independently represent hydrogen and radicals as defined for R³, or R³³ and R³⁴ together with X' represent cycloalkyl, aryl, heterocyclyl and heteroaryl

35

radicals, provided that when X' is O, R³⁴ is absent;

X' represents N, O or C(R¹⁷) wherein R¹⁷ represents hydrogen and alkyl radicals; and

5

Y, Y' and Y" independently represent O and S.

124. Compound of Claim 123 wherein Y, Y' and Y" are O.

10

125. Compound of Claim 123 wherein t is O.

126. Compound of Claim 123 wherein X' represents N and O.

15

127. Compound of Claim 126 wherein R¹ represents hydrogen and alkyl radicals having from 1 to about 4 carbon atoms, alkenyl, alkynyl, aralkyl radicals, hydroxyl radicals, and radicals represented by the formula -CH₂C(O)R" wherein R" represents R³⁸, -NR³⁸R³⁹ and -OR³⁸ wherein R³⁸ and R³⁹ independently represent hydrogen and alkyl radicals having from 1 to about 4 carbon atoms.

20

128. Compound of Claim 123 wherein R¹ represents hydrogen, methyl, ethyl, benzyl, phenylpropyl, propargyl, hydroxyl and radicals represented by the formula

25

CH₂C(O)R" wherein R" represents -CH₃, NH₂ and -OH.

129. Compound of Claim 125 wherein R¹ and R³¹ are both hydrogen and R³⁰ and R³² are both methyl.

30

130. Compound of Claim 125 wherein R³⁰ is hydrogen and R¹, R³¹ and R³² are all methyl.

35

131. Compound of Claim 125 wherein R³⁰, R³¹ and R³² are hydrogen and R¹ is methyl.

132. Compound of Claim 125 wherein R¹ and R³¹ are both hydrogen and R³⁰ and R³² together with the carbon atoms to which they are attached form a three to six-membered cycloalkyl radical.

133. Compound of Claim 125 wherein X' is O and R³³ is hydrogen or an alkyl radical.

134. Compound of Claim 125 wherein X' is O, R³⁴ is absent and R³³ represents an aralkyl radical.

135. Compound of Claim 125 wherein R² represents alkyl, cycloalkylalkyl and aralkyl radicals, which radicals are optionally substituted with halogen radicals and radicals represented by the formula -OR⁹ and -SR⁹ wherein R⁹ represents hydrogen and alkyl radicals.

136. Compound of Claim 125 wherein R² represents alkyl, cycloalkylalkyl and aralkyl radicals.

137. Compound of Claim 125 wherein R² represents aralkyl radicals.

138. Compound of Claim 125 wherein R² represents CH₃SCH₂CH₂-, iso-butyl, n-butyl, benzyl, 2-naphthylmethyl and cyclohexylmethyl radicals.

139. Compound of Claim 125 wherein R² represents an n-butyl and iso-butyl radicals.

140. Compound of Claim 125 wherein R² represents benzyl, 4-fluorobenzyl, and 2-naphthylmethyl radicals.

141. Compound of Claim 125 wherein R² represents a cyclohexylmethyl radical.

142. Compound of Claim 125 wherein R³, R⁴ and R⁵ independently represent alkyl, alkenyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, aralkyl and heteroaralkyl radicals.

143. Compound of Claim 125 wherein R⁵ represents hydrogen.

10

144. Compound of Claim 142 wherein R³ and R⁴ independently represent alkyl and alkenyl radicals.

145. Compound of Claim 143 wherein R³ and R⁴ independently represent alkyl and hydroxyalkyl radicals.

15

146. Compound of Claim 143 wherein R³ and R⁴ independently represent alkyl, cycloalkyl and cycloalkylalkyl radicals.

20

147. Compound of Claim 143 wherein R³ and R⁴ independently represent alkyl, heterocycloalkyl and heterocycloalkylalkyl radicals.

25

148. Compound of Claim 143 wherein R³ and R⁴ independently represent alkyl, aryl and aralkyl radicals.

149. Compound of Claim 143 wherein R³ and R⁴ independently represent alkyl, cycloalkyl, cycloalkylalkyl, heterocycloalkyl, heterocycloalkylalkyl, aryl, aralkyl, and heteroaralkyl radicals.

30

150. Compound of Claim 125 wherein R³ represents alkyl radicals having from about 2 to about 5 carbon atoms.

35

151. Compound of Claim 125 wherein R³ independently represent n-propyl, i-butyl, neo-pentyl, n-pentyl, n-hexyl, i-amyl, and n-butyl radicals.

5 152. Compound of Claim 125 wherein R³, R⁴ and R⁵ independently represent alkyl radicals having from about 2 to about 5 carbon atoms, cycloalkylalkyl radicals, aralkyl radicals, heterocycloalkylalkyl radicals and heteroaralkyl radicals.

10 153. Compound of Claim 125 wherein R³ represents benzyl, para-fluorobenzyl, para-methoxybenzyl, para-methylbenzyl, and 2-naphthylmethyl radicals and R⁴ represents t-butyl.

15 154. Compound of Claim 125 wherein R³ is cyclohexylmethyl and R⁴ is t-butyl.

20 155. Compound of Claim 125 wherein R³ is i-amyl or n-butyl and R⁴ is t-butyl.

156. Compound of Claim 125 wherein R³ is i-butyl and R⁴ is t-butyl.

25 157. Compound of Claim 125 wherein R³ is benzyl or p-fluorobenzyl and R⁴ is t-butyl.

158. Compound of Claim 125 wherein R³ is neo-pentyl and R⁴ is t-butyl.

30 159. Compound of Claim 125 wherein R⁴ represents alkyl and cycloalkyl radicals.

35 160. Compound of Claim 125 wherein R⁴ represents ethyl, i-propyl, t-butyl and 1,1-dimethylpropyl radicals.

161. Compound of Claim 125 wherein R³ represents heteroaralkyl radicals and R⁴ is t-butyl.

162. Compound of Claim 125 wherein R⁴ and R⁵ together with the nitrogen atom to which they are attached represent a 5 to 8-membered heterocyclic or heterobicyclicalkyl radical.

163. Compound of Claim 125 wherein R⁴ and R⁵ together with the nitrogen to which they are bonded represent pyrrolidinyl, piperidinyl, morpholinyl, and piperazinyl radicals.

164. A pharmaceutical composition comprising a compound of Claim 123 and a pharmaceutically acceptable carrier.

165. Method of inhibiting a retroviral protease comprising administering a protease inhibiting amount of a composition of Claim 164.

166. Method of Claim 165 wherein the retroviral protease is HIV protease.

167. Method of treating a retroviral infection comprising administering an effective amount of a composition of Claim 164.

168. Method of Claim 167 wherein the retroviral infection is an HIV infection.

169. Method for treating AIDS comprising administering an effective amount of a composition of Claim 164.

170. A compound of Claim 1 which is:

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-methylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl]propylamino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl]butylamino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](phenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-fluorophenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-methoxyphenylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3,3-dimethylpropyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

5

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-hydroxybutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

10

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](4-pyridylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

15

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-phenylethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

20

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](2-naphthylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

25

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](cyclohexylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

30

Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(4-fluorophenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

35

- Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(cyclohexylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;
- 5 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(butyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;
- 10 Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-;
- 15 Propanamide, N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(4-fluorophenylmethyl)propyl]-2-methyl-3-(methylsulfonyl)-, [1S-[1R*(R*), 2S*]]-;
- 20 Propanamide, 3-(acetylamino)-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-;
- 25 Propanamide, 3-(acetylamino)-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(4-fluorophenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-;
- 30 Propanamide, 3-(acetylamino)-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(cyclohexylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-;

Propanamide, 3-(phenoxyacetyl amino)-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-;

5

Propanamide, 3-(2-naphthyloxyacetyl amino)-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-;

10

Propanamide, 3-(4-methoxybenzyloxy carbonyl) amino-N-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2-methyl-, [1S-[1R*(S*),2S*]]-;

15

2,6,10,12-tetraazatetradecanamide, 8-hydroxy-N,4,13,13-tetramethyl-10-(3-methylbutyl)-5,11-dioxo-7-(phenylmethyl)-, [4R-(4R*,7S*,8R*)]]-;

20

2,6,10,12-tetraazatetradecanamide, 8-hydroxy-N,4,13,13-tetramethyl-10-(3-methylbutyl)-5,11-dioxo-7-(4-fluorophenylmethyl)-, [4R-(4R*,7S*,8R*)]]-;

25

Butanediamide, N¹-[2-hydroxy-3-[(phenylmethyl)(1-pyrrolidinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

30

Butanediamide, N¹-[2-hydroxy-3-[(4-fluorophenylmethyl)(1-pyrrolidinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

35

Butanediamide, N¹-[2-hydroxy-3-[(3-methylbutyl)(4-morpholinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

Butanediamide, N¹-[2-hydroxy-3-[(4-fluorophenylmethyl)(4-morpholinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

- 5 Butanediamide, N¹-[2-hydroxy-3-[(4-fluorophenylmethyl)(1-piperidinylcarbonyl)amino]-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-;

- 10 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-;

- 15 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(4-fluorophenylmethyl)propyl]-2,3,3-trimethyl-, [1S-[1R*(2S*),2S*]]-;

- 20 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-2,3-dimethyl-, [1S-[1R*(2S*,3R*),2S*]]-;

- 25 Butanediamide, N¹-[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(4-fluorophenylmethyl)propyl]-2,3-dimethyl-, [1S-[1R*(2S*,3R*),2S*]]-;

- 30 Butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]amino]-2,2,3-trimethyl-4-oxo-, [1S-[1R*(3S*),2S*]]-;

- 35 Butanoic acid, 4-[[3-[[[(1,1-dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-hydroxy-1-(4-fluorophenylmethyl)propyl]amino]-2,2,3-trimethyl-4-oxo-, [1S-[1R*(3S*),2S*]]-;

Butaneamide,

2-(acetylamino)-N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
5 hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-
[1R*(R*),2S*]]-;

Butaneamide,

2-(acetylamino)-N-[3-[[[(1,1-
10 dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
hydroxy-1-(4-fluorophenylmethyl)propyl]-3,3-dimethyl-, [1S-
[1R*(R*),2S*]]-;

Butaneamide,

15 2-(acetylamino)-N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl](4-fluorophenylmethyl)amino]-
2-hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-
[1R*(R*),2S*]]-;

20 Butaneamide,

2-[(2,2-dimethylaminoacetyl)amino]-N-[3-[[[(1,1-
dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
hydroxy-1-(phenylmethyl)propyl]-3,3-dimethyl-, [1S-
[1R*(R*),2S*]]-.

25

or;

Butanediamide, N¹-[3-[[[(1,1-

dimethylethyl)amino]carbonyl](3-methylbutyl)amino]-2-
30 hydroxy-1-(phenylmethyl)propyl]-3-[(2-
quinolinylcarbonyl)amino]-, [1S-[1R*(R*),2S*]]-.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 93/04806

I. CLASSIFICATION OF SUBJECT MATTER: (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5	C07C275/24; C07C275/22;	A61K31/17; A61K31/47
		C07D215/48;
		C07C317/44
II. FIELDS SEARCHED:		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07C	
Documentation Searched other than Minimum Documentation to the extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	EP,A,0 264 795 (MERCK PATENT GMBH) 27 April 1988 cited in the application see claims ----	1-68
Y	EP,A,0 374 097 (CIBA-GEIGY) 20 June 1990 see page 2, line 25 - line 20 see page 7, line 20 - line 34 ----	1-68
A	GB,A,2 184 730 (E. R. SQUIBB & SONS INC.) 1 July 1987 cited in the application see claims; examples ----	78-117
A	US,H,725 (E. M. GORDON) 2 January 1990 cited in the application see claims; examples ----	1-170
A	US,H,725 (E. M. GORDON) 2 January 1990 cited in the application see claims; examples ----	1-170
	-/--	
¹⁰ Special categories of cited documents : ¹⁰ "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "A" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 13 AUGUST 1993		Date of Mailing of this International Search Report - 1. 09. 93
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer SEUFERT G.H.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	GB,A,2 200 115 (SANDOZ) 27 July 1988 cited in the application see claims 1,2	1-170
P,X	WO,A,9 208 701 (MONSANTO CO) 29 May 1992 see claims; examples; tables	1-67, 78-117, 123-164, 170

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 93/04806

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 68-77, 118-122, 165-169 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2. ☒ Claims Nos.: 1-169 (searched incompletely)
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
The vast number of theoretically conceivable compounds resulting from the combination of all claimed substituents, as well as the use of rather vague terms like heteroaxyl, heterocyclyl preclude a comprehensive search. For economical reasons the search has been limited to the examples.
(Guidelines exam. Part B; Chapt. III; 3.6, 3.7)
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9304806
SA 74473

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 13/08/9

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0264795	27-04-88	DE-A- 3635907 AU-A- 7982387 JP-A- 63112548 ZA-A- 8707950	28-04-88 28-04-88 17-05-88 26-04-88
EP-A-0374097	20-06-90	AU-A- 4678589 CA-A- 2005337 JP-A- 2212434	21-06-90 15-06-90 23-08-90
GB-A-2184730	01-07-87	CA-A- 1297631 DE-A- 3643977 FR-A- 2592048 JP-A- 62164658 US-A- 4757050	17-03-92 02-07-87 26-06-87 21-07-87 12-07-88
US-H-725	02-01-90	None	
GB-A-2200115	27-07-88	AU-A- 1037588 BE-A- 1002212 CH-A- 676988 DE-A- 3800591 FR-A- 2609716 JP-A- 1019053 LU-A- 87110 NL-A- 8800100 SE-A- 8800169	01-09-88 16-10-90 28-03-91 04-08-88 22-07-88 23-01-89 23-08-88 16-08-88 22-07-88
WO-A-9208701	29-05-92	AU-A- 9053191 AU-A- 9085191 AU-A- 9092591 AU-A- 9125191 AU-A- 9133291 EP-A- 0554400 WO-A- 9208698 WO-A- 9208699 WO-A- 9208700 WO-A- 9208688	11-06-92 11-06-92 11-06-92 11-06-92 11-06-92 11-08-93 29-05-92 29-05-92 29-05-92 29-05-92